



# Department for Environment Food & Rural Affairs



## Monthly publication of National Statistics on the Incidence of Tuberculosis (TB) in Cattle to end June 2014 for Great Britain

These statistics were released today, Wednesday 10 September 2014 at 09:30, with the next notice to be updated on Wednesday 15 October 2014 at 09:30.

These statistics are obtained from the Animal Health and Veterinary Laboratories Agency (AHVLA) work management IT support system (Sam), used for the administration of TB testing in GB. They are a snapshot of the position on the date on which the data were extracted. These statistics may be subject to regular revision until all test results are available. In particular figures from 2012 onwards will be subject to further revision as test and incident records are completed.

### The key points relating to June 2014 are:-


- Short term changes in these statistics should be considered in the context of long term trends. The charts and tables in this statistical notice give the latest indication of how the trend in bovine TB incidence has changed since 1996.
- The provisional incidence rate for January to June 2014 is 3.6% compared to 4.2% for January to June 2013. However, care needs to be taken not to read too much into short term figures, especially as this figure includes a number of unclassified incidents. As such, the incidence rates are subject to further revisions as more tests and their results for the period are input.
- The number of new herd incidents during the period January to June 2014 was 2,398 compared to 2,535 for January to June 2013. The number of tests on officially TB free herds was 42,819 during January to June 2014, compared to 39,439 during January to June 2013.
- The number of cattle compulsorily slaughtered as reactors or direct contacts was 17,063 during January to June 2014, compared to 17,282 during January to June 2013.

---


Enquiries to :- [tbstatistics@defra.gsi.gov.uk](mailto:tbstatistics@defra.gsi.gov.uk)  
Media Enquiries to :- 0207 238 6007 (Press Office)

**A National Statistics publication.** National Statistics are produced to high professional standards. They undergo regular quality assurance reviews to ensure that they meet customer needs. They are produced free from any political interference. For general enquiries about National Statistics, contact the National Statistics Public Enquiry Service: tel. 0845 601 3034 email [info@statistics.gov.uk](mailto:info@statistics.gov.uk). You can find National Statistics on the internet at [www.statistics.gov.uk](http://www.statistics.gov.uk).

**Figure 1: Herd incidence of bovine TB (bTB) in GB: number of new incidents of bTB leading to the withdrawal of officially TB free (OTF) herd status, as a percentage of tests carried out in OTF herds each month since 2003:**



**Figure 2: Herd incidence of bovine TB (bTB) in GB: number of new incidents of bTB leading to the withdrawal of officially TB free (OTF) herd status, as a percentage of tests carried out in OTF herds each month since 1996:**



There has been an overall long-term upward trend in the incidence of TB in cattle herds in GB since the beginning of the time series presented in this statistical release although the incidence rate is lower now than it was at its peak in 2008.

An unusual spike in the incidence rate was observed during most of 2001. This was an anomaly caused by the suspension of TB testing during the foot and mouth disease (FMD) outbreak of February to October 2001. In the meantime, new bTB breakdowns continued to be detected on farms through routine post-mortem meat inspection of cattle carcasses in abattoirs. Following the gradual resumption of herd testing after the 2001 FMD outbreak, a three-year cycle in the bTB herd incidence rate was observed (as shown in two figures above and in Table 1), with peaks in early 2005, early 2008 and in 2011 and troughs in 2006 and 2009.

There is no clear explanation for this pattern, because bTB is a multi-factorial and chronic disease with a complex epidemiology and reservoirs of infection in cattle and wildlife. As a result of surveillance and testing changes, there has been no stable time series until recently. For example there have been different herd testing frequencies in each parish over time, ranging from annual to four-yearly and changing every year until those frequencies were unified in Wales in 2010 (annual) and in England in 2013 (annual and four-yearly).

There are at least three possible explanations:

1. The smoothed trend represents true seasonal changes in the transmission risk and prevalence of infection in wildlife and cattle populations. However there is no strong evidence to support this.
2. After FMD higher risk herds were tested every 3 to 4 years and could have contributed to a cyclically higher incidence rate. However breakdowns in the 4-yearly (and formerly 3-yearly) testing areas represent a small and decreasing proportion of the breakdowns in any given year.
3. When testing resumed in 2002 following the 2001 FMD outbreak, high-risk herds may have been identified then put under restrictions and control tested for a period (when they cannot generate a new incident). Once the bTB incident has been resolved and OTF herd status is restored, the herd becomes susceptible to a new incident as it undergoes post-breakdown surveillance tests at 6 and 18 months after regaining OTF status.

The charts published in this statistical notice, together with the equivalent figures from January 1996 onwards, are also available in spreadsheet format on the Defra web site at :  
<https://www.gov.uk/government/publications/incidence-of-tuberculosis-tb-in-cattle-in-great-britain>. Visit the third link (MS Excel spreadsheet) entitled “Incidence of TB in cattle in Great Britain - GB dataset”.

**TABLE 1: TB INCIDENTS IN GREAT BRITAIN - HERDS**

		Number of cattle herds registered on Sam	Total tests on herds	Herds not Officially TB free due to a bovine TB incident (non-OTF Herds)	Tests on officially TB free herds (OTF)	New herd incidents (NHI)	NHI of which: officially TB free herd status withdrawn (OTFW)	Number of OTFW incidents as a percentage of tests on officially TB free herds
		(1)	(2)	(3)	(4)	(5)	(6)	(7)
1996		na	36,314	1,589	34,812	1,075	490	1.4%
1997		na	34,065	1,632	32,295	1,195	540	1.7%
1998		na	37,046	2,077	34,502	1,514	787	2.3%
1999		na	41,365	2,374	38,338	1,661	967	2.5%
2000		na	40,669	2,482	37,184	1,738	1,135	3.1%
2001	*	na	13,187	1,697	11,118	802	571	5.2%
2002	**	na	49,709	4,167	43,641	3,323	2,042	4.7%
2003		na	56,208	5,460	47,568	3,214	1,789	3.8%
2004		na	56,836	5,220	49,027	3,341	1,934	4.0%
2005		na	55,887	5,669	46,725	3,665	2,308	4.9%
2006		na	64,457	5,859	56,051	3,530	2,303	4.1%
2007		na	64,145	6,582	54,856	4,188	2,546	4.7%
2008		na	66,432	7,935	54,854	5,011	3,093	5.6%
2009		na	72,205	8,386	58,894	4,599	2,847	4.9%
2010		83,636	74,474	7,964	61,587	4,723	3,013	4.9%
2011		80,426	76,659	8,243	62,489	4,912	3,112	5.2%
2012	(p)	79,251	88,587	8,958	73,662	5,151	3,469	4.8%
2013	(p)	79,287	86,846	9,123	72,181	4,821	3,255	4.5%
2014	(p)	78,834	50,405	6,312	42,819	2,398	1,538	3.6%
2012	Jan	(p)	80,383	8,195	4,292	6,996	485	326 - 333
	Feb	(p)	80,331	9,059	4,406	7,699	464	315 - 326
	Mar	(p)	80,203	11,719	4,605	10,284	592	400 - 408
	Apr	(p)	80,174	7,891	4,649	6,534	405	248 - 254
	May	(p)	79,986	7,305	4,692	5,992	448	261 - 269
	Jun	(p)	79,986	5,428	4,557	4,178	325	200 - 203
	Jul	(p)	79,857	5,068	4,482	3,865	315	217 - 219
	Aug	(p)	79,717	5,422	4,440	4,243	353	244 - 249
	Sep	(p)	79,604	5,882	4,392	4,743	367	264 - 270
	Oct	(p)	79,496	6,816	4,492	5,747	444	328 - 333
	Nov	(p)	79,350	9,066	4,618	7,682	558	392 - 400
	Dec	(p)	79,251	6,736	4,624	5,699	395	274 - 276
2013	Jan	(p)	79,187	8,753	4,800	7,342	498	312 - 321
	Feb	(p)	79,188	9,004	4,813	7,561	398	280 - 283
	Mar	(p)	79,241	9,256	4,834	8,011	489	317 - 323
	Apr	(p)	79,354	8,082	4,816	6,608	423	286 - 292
	May	(p)	79,469	7,354	4,647	5,970	402	265 - 270
	Jun	(p)	79,553	5,088	4,434	3,947	325	193 - 199
	Jul	(p)	79,558	5,081	4,292	3,860	278	175 - 179
	Aug	(p)	79,538	5,551	4,160	4,414	328	224 - 227
	Sep	(p)	79,501	5,982	4,123	4,913	350	252 - 253
	Oct	(p)	79,511	7,273	4,188	6,207	472	340 - 344
	Nov	(p)	79,426	8,583	4,264	7,444	503	360 - 363
	Dec	(p)	79,287	6,839	4,262	5,904	355	251 - 255
2014	Jan	(p)	79,111	9,679	4,448	8,120	534	368 - 369
	Feb	(p)	79,052	9,046	4,448	7,892	429	284 - 285
	Mar	(p)	78,756	9,580	4,474	8,447	445	295 - 301
	Apr	(p)	78,798	8,080	4,408	6,826	360	216 - 219
	May	(p)	78,837	8,303	4,266	6,944	374	222 - 227
	Jun	(p)	78,834	5,717	4,041	4,590	256	153 - 155

**Notes:-** The data are a snapshot extracted from Sam. Data for 2012 onwards will remain provisional and subject to revision each month until all culture results are available and final data validation has been carried out. The herd incidence rates for the latest months are given as a range because a number of incidents are still unclassified, so data for these months should be treated as provisional results. TB incidents remain unclassified if at the end of the period covered by this notice they had not been designated OTFW, but were still ongoing and the herd could have its OTF status withdrawn if further testing revealed one or more animals with post-mortem evidence of TB.

- (1) The number of herds registered on the AHVLA's Sam (computer) system – monthly figures are not available (na) before December 2010.
- (2) Herds for which tuberculin skin testing is carried out on at least one animal during the period shown. Does not include gamma tests. (same as column 1 in Table 2).
- (3) Herds that had lost their OTF status at some time during the period shown due to a TB incident.
- (4) Any test carried out in an OTF herd during the period shown. Does not include gamma tests.
- (5) Herds which were previously OTF but either had cattle that reacted to a tuberculin test or had a tuberculous animal disclosed by routine meat inspection at slaughter, during the period shown.
- (6) New herd incidents (column 4) where OTF status was withdrawn from the herd.
- (7) Column 5 as a percentage of column 3.
- \* Data for 2001 are not comparable with other years. During the outbreak of Foot and Mouth Disease, TB testing was significantly reduced and necessarily targeted to areas of higher risk.
- \*\* Data for 2002 are not comparable with other years. Testing resources were concentrated on herds overdue their tests (because of the backlog caused by the Foot and Mouth Disease outbreak).
- (p) provisional

**TABLE 2: TB INCIDENTS IN GREAT BRITAIN - ANIMALS**

	Total tests on herds (1)	Total cattle tests (2)	Cattle compulsorily slaughtered as reactors or contacts:		
			Total (3)	Reactors (4)	Direct contacts (5)
1996	36,314	2,249,891	3,776	3,151	625
1997	34,065	2,170,630	3,384	3,017	367
1998	37,046	2,447,848	5,685	4,782	903
1999	41,365	2,825,177	6,754	5,794	960
2000	40,669	2,931,658	8,123	6,877	1,246
2001	* 13,187	1,181,861	6,156	5,200	956
2002	** 49,709	3,961,145	22,072	19,191	2,881
2003	56,208	4,474,526	23,972	20,798	3,174
2004	56,836	4,604,721	22,214	19,636	2,578
2005	55,887	4,811,699	29,231	25,627	3,604
2006	64,457	5,417,573	22,062	20,090	1,972
2007	64,145	5,753,244	26,882	25,330	1,552
2008	66,432	6,178,789	39,007	36,968	2,039
2009	72,205	6,840,568	37,979	36,739	1,240
2010	74,474	7,447,653	31,949	31,277	672
2011	76,659	7,587,837	34,238	33,453	785
2012	(p) 88,587	8,031,649	37,735	37,050	685
2013	(p) 86,846	8,393,303	32,620	31,723	897
2014	(p) 50,405	4,805,001	17,063	16,482	581
2012	Jan (p) 8,195	720,548	2,580	2,537	43
	Feb (p) 9,059	781,628	3,769	3,714	55
	Mar (p) 11,719	990,671	3,124	3,090	34
	Apr (p) 7,891	722,223	2,805	2,774	31
	May (p) 7,305	580,183	3,467	3,414	53
	Jun (p) 5,428	491,658	2,527	2,461	66
	Jul (p) 5,068	477,630	3,311	3,244	67
	Aug (p) 5,422	487,429	2,989	2,864	125
	Sep (p) 5,882	547,355	2,637	2,619	18
	Oct (p) 6,816	656,406	3,935	3,874	61
	Nov (p) 9,066	932,913	3,753	3,681	72
	Dec (p) 6,736	643,005	2,838	2,778	60
2013	Jan (p) 8,753	771,196	3,200	3,141	59
	Feb (p) 9,004	814,718	3,105	2,997	108
	Mar (p) 9,256	854,194	2,973	2,860	113
	Apr (p) 8,082	825,434	2,726	2,690	36
	May (p) 7,354	664,875	3,243	2,962	281
	Jun (p) 5,088	482,595	2,035	2,004	31
	Jul (p) 5,081	512,932	2,706	2,651	55
	Aug (p) 5,551	537,364	2,523	2,470	53
	Sep (p) 5,982	575,842	2,105	2,067	38
	Oct (p) 7,273	722,179	2,855	2,811	44
	Nov (p) 8,583	926,540	2,750	2,715	35
	Dec (p) 6,839	705,434	2,399	2,355	44
2014	Jan (p) 9,679	998,876	2,923	2,883	40
	Feb (p) 9,046	863,595	3,001	2,868	133
	Mar (p) 9,580	880,890	2,896	2,847	49
	Apr (p) 8,080	798,576	2,866	2,747	119
	May (p) 8,303	717,401	2,724	2,690	34
	Jun (p) 5,717	545,663	2,653	2,447	206

**Notes:** The data are a snapshot extracted from Sam. Data for 2012 onwards will remain provisional and subject to revision each month until all culture results are available and final data validation has been carried out.

- (1) Herds for which tuberculin skin testing is carried out on at least one animal during the period shown. Does not include gamma tests. (same as column 2 in Table 1).
  - (2) Count of the number of tests on cattle. An individual animal could be tested more than once in each time period.
  - (3) Animals compulsorily slaughtered because they reacted to the tuberculin skin test or because they were considered to be direct contacts (see below). Not all of these animals showed evidence of *Mycobacterium bovis* infection at post-mortem examination.
  - (4) An animal which was compulsorily slaughtered because it responded to the tuberculin skin test in a way that was consistent with it being infected with *Mycobacterium bovis*.
  - (5) An animal in an OTFW incident that, although not a test reactor, was considered to have been exposed to *Mycobacterium bovis* and compulsorily slaughtered.
- \* Data for 2001 are not comparable with other years. During the outbreak of Foot and Mouth Disease, TB testing was significantly reduced and necessarily targeted to areas of higher risk.
- \*\* Data for 2002 are not comparable with other years. Testing resources were concentrated on herds overdue their tests (because of the backlog caused by the Foot and Mouth Disease outbreak).
- (p) provisional

## What is bovine tuberculosis?

Bovine tuberculosis (bTB) is a chronic infectious disease of cattle<sup>1</sup>. The risk bTB poses to human health is low, largely due to milk pasteurisation. The disease is detected either on farms (through mandatory skin tests<sup>2</sup> of cattle herds for bTB at regular intervals) and at abattoirs (through post-mortem meat inspection of cattle carcasses).

## What are the impacts of bTB?

Bovine TB presents serious challenges to the food and farming industries and has economic and social impacts. The economic costs of a bTB breakdown<sup>3</sup> are shared by farmers and government; in 2012 the estimated average cost of a confirmed herd breakdown in high risk areas of England was £14,000 to farmers and £20,000 to government<sup>4</sup>. Costs are incurred for a number of reasons:

- Cattle which are found (or are highly likely) to have bTB are slaughtered. This loses the farmer the value of the animal and its output. Government pays farmers compensation for slaughtered animals which is based on the market value of cattle.
- There are costs associated with testing animals for bTB. Farmers incur costs from gathering animals together, such as paying workers for their time, and government pays the vets' fees for carrying out tests on the herd (and in the event of a breakdown on herds in neighbouring farms).
- When an animal in a herd tests positive for the disease, the whole herd is put under movement restrictions until all the remaining animals are tested repeatedly with negative results. This presents costs to farmers, for example because they are unable to move their cattle to market or buy in replacements for animals that are slaughtered.

Other impacts of high bTB levels can include:

- Restrictions on trade in cattle within Europe<sup>5</sup>
- Significant stress amongst farmers, their families and local communities<sup>6</sup>
- The infection spilling over to domestic and wild animals<sup>7</sup>.

## Why monitor statistics about bTB?

**Legal requirements:** EU Member States are legally required to have accelerated bTB eradication plans in place in order to achieve officially TB free (OTF) status<sup>8</sup>. Defra and Welsh Government policy is to achieve OTF status for the whole country by 2038, while Scotland achieved OTF status in September 2009. bTB statistics are used in England and Wales to measure progress towards this

<sup>1</sup> bTB is caused by the bacterium *Mycobacterium bovis* (*M. bovis*). Cattle are the natural host of the bacterium, but many other species, including wildlife such as badgers and (less commonly) deer, are also susceptible to *M. bovis*, can develop TB and transmit the infection to other species.

<sup>2</sup> the tuberculin skin test: if tuberculin (a purified sterile cocktail of proteins derived from *M. bovis* cultures) is injected into the skin of an animal infected with *M. bovis*, this will cause a localised allergic reaction characterised by temporary swelling of the skin, which is measured 72hrs after the injection. The principle is very similar to the skin tests for TB in humans.

<sup>3</sup> A breakdown is the term used to describe the occurrence in a herd of at least one animal with a positive reaction to the skin test, or the identification of *M. bovis* in an animal with TB lesions detected at routine slaughter. The affected herd is then placed under restrictions and loses its Officially TB Free (OTF) status.

<sup>4</sup> Economic analysis based on [research report SE3112 for Defra, 2004](#)

<sup>5</sup> Because the disease undermines the effective operation of the single market – see the [EU Animal Health Strategy](#)

<sup>6</sup> See for example [research report SE3120 for Defra, 2008](#)

<sup>7</sup> For example Broughan, J. M., Downs, S. H., Crawshaw, T. R., Upton, P. A., Brewer, J. & Clifto-Hadley, R. S. (2013) *Mycobacterium bovis* infections in domesticated non-bovine mammalian species. Part 1: review of epidemiology and laboratory submissions in Great Britain 2004-2010. *Veterinary Journal* **198**, 346-35. See also <http://www.defra.gov.uk/animal-diseases/a-z/bovine-tb/animal-keepers/other-species/>

<sup>8</sup> "OTF Status" takes its meaning from European law: for a region or Member State of the EU to be considered to be OTF the annual incidence of herds with confirmed *M. bovis* infection must not have exceeded 0.1% and at least 99.9% of the herds within it must have been free from bTB at the end of the year for at least six consecutive years.

target, and to support the annual case for Scotland to retain its OTF status, as the qualification is based on herd incidence.

**Monitoring policy effectiveness:** Statistics on the incidence of bTB in cattle herds and the number of cattle slaughtered as a result of bTB are used by policymakers to monitor the spread and concentration of the disease and to inform decisions around the potential approaches to controlling it. Existing controls include routine testing in cattle based on the disease incidence (or risk) in a given area, restricting movements of cattle from herds where an animal has tested positive for the disease and addressing the problem of disease spread through wildlife (principally badgers).

### Factors affecting statistics on incidence of bTB in cattle herds

Variation in the monthly statistics can occur for a number of reasons, including:

- **Disease:** an increase in the trend can be the result of a higher proportion of herds experiencing a breakdown because of an increase in the underlying incidence of bTB.
- **Surveillance policy** (including the frequency of testing): Cattle herds in high risk areas<sup>9</sup> are tested annually and cattle herds in low risk areas are usually tested every four years. If cattle herds in a low prevalence region are tested more frequently than every four years, the increase in the number of bTB tests will not necessarily be followed by a similar increase in the detection of infected cattle and so this may result in a decline in the incidence rate.
- **Seasonality:** more animals are tested when they are housed, during winter months, compared with when they are grazing outdoors in summer months. This is simply because it is easier to gather and test the cattle when they are already contained within a building. The blue trend line in Figures 1 and 2 account for this by presenting seasonally adjusted data.
- Number of **testing days** in a given month: tests tend to be carried out at the beginning of the working week and the results collected and entered into the data system towards the end of the week. Months containing five Fridays may therefore have more positive test results than months containing four.

An extreme example of the impact of testing on the incidence rate can be seen in the statistics for 2001, when bTB testing was significantly reduced for most of the year due to the outbreak of Foot and Mouth Disease but new bTB breakdowns continued to be detected through disease surveillance in abattoirs. This led to an unusually high incidence rate for 2001 and 2002, when effectively two years' worth of breakdowns were identified in one year when the normal testing regime resumed.

### Surveillance policy in GB

These statistics are presented for GB, but the bTB surveillance and control policy – including how frequently animals are tested for bTB – varies between England, Wales and Scotland and has changed over time.

#### Timeline:

- **1990s:** most herds in GB tested every four years and background testing intervals determined on a parish basis. Herds in parishes with a high incidence of bTB breakdowns (in the South West of England and in parts of Wales) are tested on an annual or biennial basis, with a smaller number of three-yearly testing herds.
- **2004 to 2010:** the proportion of parishes and herds in England and Wales with annual testing increases gradually as the disease spread, with a corresponding decrease in the proportion of parishes with four-yearly testing.

---

<sup>9</sup> South West, West Midlands and East Sussex, where the majority of TB cases are found and where the prevalence (probability) of TB-infected cattle and badgers is relatively high.

- **October 2009:** the European Commission designates Scotland as an officially bTB free region of the UK.
- **January 2010:** In England, a core annual testing area is established, spanning entire counties in the South West and West Midlands (the 'high risk area') and surrounded by a 'buffer' of two-yearly testing parishes. Most of the rest of England remains on background four-year testing. The Welsh Government puts all cattle herds in Wales on annual bTB testing (with herds in the small Intensive Action Area of West Wales put on 6-monthly bTB testing).
- **2011 and 2012:** further expansion of the annual testing area in England to the east and north.
- **January 2013:** herd testing intervals are determined on a county basis and England is split into annual testing and four-yearly testing counties. Annual testing of herds is extended to all the counties at the edge of the high risk area (more detail below). Three- and two-yearly testing is abolished.

### **Current differences in surveillance policy in GB**

- **England** is divided into two cattle bTB testing frequency areas that broadly reflect the geographically clustered nature of the disease. The majority of bTB cases are found in counties of the South West, West Midlands and East Sussex. These herds are tested for bTB annually and represent nearly 60% of all herds in England. In the rest of England most herds are tested every four years. Herds that have a high risk of contracting bTB or present a potential public health risk (e.g. producer-retailers of unpasteurised milk) are tested annually regardless of their location.
- All herds in **Wales** are tested annually.
- **Scotland** has in place a risk-based routine herd testing policy. This targets testing at higher risk herds. Around 35 per cent of herds are considered low risk herds and are exempt from routine testing. Herds that are not exempt tested are every four years.

More information on bovine TB can be found at:

England :-

<https://www.gov.uk/government/policies/reducing-bovine-tuberculosis>

Wales :-

<http://wales.gov.uk/topics/environmentcountryside/ahw/disease/bovinetuberculosis>

Scotland :-

<http://www.scotland.gov.uk/Topics/farmingrural/Agriculture/animal-welfare/Diseases/disease/tuberculosis>

### **Methodology**

For a description of the data sources and methodology used in the calculation of the TB statistics, together with notes on data revisions policy etc. Refer to the Annex document at :-

[https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/305850/bovinetb-annex-24apr14.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/305850/bovinetb-annex-24apr14.pdf)

### **Further Information**

This statistical notice and a wide range of other statistics are available on the internet at -

<https://www.gov.uk/government/organisations/department-for-environment-food-rural-affairs/about/statistics>