



UK Health
Security
Agency

Ebola: Information for humanitarian aid and other workers intending to work in Ebola-affected countries in Africa

Background

Ebola virus is introduced into the human population through close contact with the blood, secretions, organs or other bodily fluids of infected animals. Once a person comes into contact with an animal that is infected with the virus, it can spread from human to human. Infection occurs from direct contact (through broken skin or mucous membranes) with the blood, or other body fluids (vomit, stool, urine, saliva, semen) of infected symptomatic people. Semen can contain virus for at least 3 months after apparent recovery from the illness.

Infection can also occur if broken skin or mucous membranes of a healthy person come into contact with environments that have become contaminated with an Ebola patient's infectious fluids such as soiled clothing, bed linen, or used needles.

The incubation period is from 2 to 21 days. The illness usually begins abruptly with fever, headache, joint and muscle pain, sore throat and intense weakness. Stomach pains, diarrhoea and vomiting may occur. Some individuals may develop a rash, red eyes, hiccups, and internal or external bleeding. In severe cases patients develop failure of the liver and kidneys.

Anyone who has close contact with a person infected with the virus, or someone who handles samples from patients is at risk. This includes hospital staff, family members and laboratory workers, all of whom are at risk if they do not use proper barrier nursing techniques. These precautions aim to provide coverage of all body sites that could be exposed to virus and include the correct use of protective gowns, gloves, and masks, in addition to wearing eye protection (such as goggles) or a face shield.

Ebola virus is not spread through routine, social contact (such as shaking hands) with asymptomatic individuals. The likelihood of contracting Ebola virus disease is considered very low unless there has been a specific exposure, such as direct contact with body fluids (for example, blood, faeces, vomit) of infected people with symptoms or objects that have been contaminated with body fluids.

Fevers in persons who have travelled to Ebola transmission areas are more likely to be caused by common infections, such as gastroenteritis, malaria or typhoid fever, but such persons should be evaluated by a healthcare provider to be sure.

A history of prior Ebola vaccination, if vaccine was available and administered, does not exclude the need to be assessed for Ebola virus disease if a person has potential exposure risks and develops symptoms within the 21-day incubation period.

If you travel to an area affected by an Ebola outbreak, the following precautions are recommended.

Before you leave

Before you leave:

- your organisation should be able to provide you with all the relevant personal protective equipment (PPE) for you to safely carry out your duties but check with them what will be available. In addition, a personal supply of surgical masks and disposable gloves may also be useful, but this should be discussed with your organisation
- discuss with your organisation about what you should take with you, this will usually involve assembling a travel health kit containing basic first aid essentials, such as sunscreen, insect repellent containing DEET, painkillers, anti-diarrhoea tablets, a thermometer, household disinfectant, and alcohol-based hand rubs for hand hygiene. Be aware that hand rubs will not provide protection from any infection if hands are visibly soiled; in which case washing with soap and water is required
- seek information about the current Ebola outbreak. This will be an evolving situation so remain aware of up-to-date, reliable information on the specific areas affected, as these may change over time (see the [Additional Information links](#))
- be sure you are up to date with all your routine immunisations, and see your healthcare provider at least 4 to 6 weeks before travel to review your immunisation status, and get advice on any additional travel-related immunisations, medications (most importantly malaria prophylaxis), or information you may need. However, even if you are travelling last minute, it is still not too late to get travel health advice. Some vaccines can be given at short notice and anti-malarial tablets can be started just before you travel. See [TravelHealthPro](#)
- if you are on prescribed medications, ensure you have a sufficient supply with you to cover your time in the field. If you have a pre-existing medical condition, you should discuss the suitability of this trip with your doctor or occupational health department
- if you have travel insurance, check your policy to establish the duration of trip covered for, whether emergency assistance is available, and whether this type of trip would be covered, or get additional insurance that covers medical evacuation in the event of illness. Check your organisation's health and safety policies and medical evacuation arrangements
- identify in-country healthcare resources in advance of your trip and discuss with your organisation

- your organisation will pre-register you with the [UKHSA's Returning Workers Scheme](#). UKHSA should be notified if there are changes to your arrangements for returning to the UK, particularly changes to travel dates or where you will arrive.
- if for any reason you have not been pre-registered with the scheme, see [below](#)
- you may wish to discuss the potential implications of your deployment, especially return-to-work arrangements (see 'after your return', below), with your regular employer in the UK; this is particularly important for those who do clinical work

While you are in an area where Ebola has been reported

Follow basic precautions, for example:

- avoid contact with ill or dead people unless wearing appropriate PPE
- avoid contact with ill or dead animals (especially primates and bats) as well as preparation and consumption of 'bushmeat' (also called wild meat; meat of terrestrial wild or feral mammals, killed for food in the tropics)
- adhere to safe sex practices including the use of barrier methods

If you are involved in medical care you should maintain the general principles of infection control at all times, including:

- careful and frequent hand washing using soap and water (or waterless alcohol-based hand rubs when soap is not available)
- wearing gloves
- proper disposal of needles and other equipment and sterilisation of non-disposable equipment
- proper disposal of body fluids and tissues from patients

In addition, you must observe strict barrier techniques when in close contact with persons suspected or known to have Ebola. For healthcare workers working with Ebola patients in African healthcare settings, deploying medical organisations are likely to have infection control and PPE protocols that their workers have to follow. Training in the correct use and removal of PPE is essential before any direct patient contact or other high risk work; this should be provided by the responsible medical organisation.

The World Health Organization (WHO)'s [Interim Infection Prevention and Control Guidance for Care of Patients with Suspected or Confirmed Filovirus Haemorrhagic Fever in Health-Care Settings, with Focus on Ebola](#), published in December 2014, provides guidance on the safe delivery of direct patient care, laboratory activities, post mortem examinations, movement and burial of human remains, cleaning and waste disposal.

The WHO website has a [web page devoted to Ebola](#).

Vital barrier techniques aim to cover all potential sites of exposure and include wearing protective gowns, gloves, masks and eye protection or face shields. It is vitally important that

these are very carefully removed after use. Individuals should refer to their associated medical organisation's protocols. If these are not available, then refer to the [detailed WHO advice](#).

In case of infection

If you think you have been exposed to Ebola while in the affected area:

- if you do not have any symptoms but think that you have been exposed to Ebola virus (for example, through a breach in PPE), contact your organisation immediately. They will be able to give further advice
- if you become unwell, inform your organisation, visit a healthcare provider immediately, as per your organisation's sickness management protocol, and inform them that you may have been exposed to Ebola virus. There may be a legal duty to report this and your organisation should be aware of this requirement. When travelling to seek medical attention, limit your contact with others. All other travel should be avoided
- remember that symptoms of Ebola virus disease can be non-specific, for example, chills, muscle aches, sore throat. Other potential causes of illness (such as malaria) should not be overlooked

If you are not pre-registered with the Returning Workers Scheme

Please call the [Returning Workers Scheme](#) on 020 8745 7209 (available 24 hours) and they will assist you. If you have already returned to the UK, you should contact RWS as soon as possible and within 24 hours of your arrival. If you have already been in the UK for more than 24 hours, you should still contact RWS.

After your return

On your arrival in the UK you should comply with any public health screening process for returnees, including those performed at the port of entry if they have been introduced.

Healthcare, public health and social care workers returning from an affected area will need to be assessed before they can return to work involving patient contact in the UK. This assessment will look at the type of work that you carried out in Africa and whether there were any potential exposures or issues around the use of PPE during this period. This will determine the degree of follow-up required and the work activities you can resume in the following 21 days. People are not considered infectious while asymptomatic but, to maintain high standards of patient safety within the NHS, returning workers in specific categories may have restrictions placed on the type of work they can do, until they are past the incubation period for Ebola (21 days).

Your organisation will provide your contact details and your exposure categorisation to UKHSA when you return from your work in Africa. Before you can return to work, you have a duty to

undertake the assessment (as above) and follow the advice provided. The follow-up required for each category of returning worker is dependent on the type of work undertaken.

Category 1

This may include logistics and operational support workers. There are no restrictions on the activities of a person in this category – you can return to usual activities. You will be contacted by UKHSA on your return. There is no further monitoring or reporting requirement after this contact. You will be given advice about reporting illness and accessing healthcare should you become unwell.

Category 2

This includes among others: water, sanitation and hygiene (WASH) staff (including coordinators, water engineers, hygienists and laundry staff); epidemiologists (especially in the community); health advisors (in Ebola treatment centres (ETCs) or the community); contact tracers in the community.

There will be some restrictions on what activities and work you are able to undertake in the following 21 days:

- you may return to work but, if a healthcare worker, you must not undertake exposure prone procedures for 21 days
- do not share toothbrushes or razors
- use barrier contraception or avoid unprotected sex for 21 days
- postpone any non-essential medical or dental treatment: inform healthcare provider of travel and work if any essential treatment needed

You will be contacted by UKHSA on your return. You will be required to check your temperature twice daily for 21 days after return (please see [Use of medicine](#) below), and to report any raised temperature (37.5°C or higher) or other suspicious symptoms to a designated UKHSA contact. A monitoring kit and detailed guidance will be provided by UKHSA.

Category 3

This includes healthcare workers providing patient care, morgue workers and burial teams.

There will be some restrictions on what activities and work you are able to undertake. For 21 days after returning from an Ebola-affected country the following restrictions apply:

- if a healthcare worker, no patient contact for 21 days
- postpone any non-essential medical or dental treatment: inform healthcare provider of travel and work if any essential treatment needed
- do not share toothbrushes or razors and use barrier contraception or avoid unprotected sex for 21 days

- avoid situations where you may be unable to extricate yourself quickly if you became ill (in particular crowded public places such as sports grounds)
- stay in private accommodation only, unless an alternative has been explicitly agreed between the deploying organisation and UKHSA

Travel after arrival home (while no symptoms) – any length of trip by private car within the UK is permitted, as is local travel of up to 2 hours on local commuter or public transport (principally for work). Such trips should be as short as possible. Air travel within the UK during this period is not acceptable.

You will be contacted by UKHSA on your return. You will be required to check your temperature twice daily for 21 days after return (please see [Use of medicine](#) below), and to REPORT DAILY to a designated UKHSA contact, even if you don't have a raised temperature (37.5°C or higher) or other suspicious symptoms. A monitoring kit and detailed guidance will be provided by UKHSA.

[Use of medicines such as aspirin, paracetamol or ibuprofen](#)

Certain medicines can have the effect of reducing body temperature during a fever (antipyretic). Medicine with an antipyretic, including aspirin, paracetamol or ibuprofen, can lower your body temperature for up to 8 hours. Therefore, for up to 8 hours after taking an antipyretic, thermometer measurements may show a lower temperature than would be expected due to the effect of the antipyretic.

You must contact your designated UKHSA contact if you need to take such a medicine for further advice. Please take your temperature before you take aspirin, paracetamol or ibuprofen, and if it is 37.5°C or higher then you must contact your designated UKHSA contact immediately.

A temperature of 37.5°C is always significant, whether you are taking one of these medicines or not, and must be reported urgently.

[If you become unwell within the 21-day period after return to the UK](#)

This must be reported to the designated UKHSA contact who will arrange for you to be medically assessed. If there is any delay in making contact with them, you should phone 111 reporting your recent work in an Ebola-affected area. If you are seriously ill, dial 999, and report your recent work in an Ebola-affected area.

A history of Ebola vaccination, if vaccine was available and administered before you started your work, does not exclude the need to report your illness immediately.

[Looking after yourself when you return](#)

Supporting the response in Africa will be rewarding, but it can be stressful and emotionally demanding. It is likely that you will be in contact with people who are seriously ill and their

families. It is normal to feel distress under these circumstances and your training and your colleagues will help you to deal with these challenges.

There are a number of simple things you can do while you are there and when you return home that have been shown to help people cope with difficult circumstances:

- talk to people around you: colleagues, friends and try to keep in touch with your family. Just engaging in normal activities with others can help you to cope
- do not drink too much alcohol; although it can be tempting and seem to help in the short term, drinking too much can make things worse
- look after yourself physically: do not overdo it; take breaks; remember to eat regularly and properly; try to get enough sleep; gentle exercise can help
- if you are struggling to cope then you should tell your supervisor or your GP who can help you to find further support. Remember that if you look after yourself and your own needs, you will be in a better position to help others

Additional information

More information can be found on the following websites:

- [Health recommendations for travel to Africa](#)
- [Information about current outbreaks](#)
- [More information about Ebola virus disease](#)
- [Information about other viral haemorrhagic fevers](#)
- [Viral haemorrhagic fever: ACDP algorithm and guidance on management of patients](#)

About the UK Health Security Agency

The UK Health Security Agency is an executive agency, sponsored by the [Department of Health and Social Care](#).

© Crown copyright 2021
Version 8

Published: November 2021
UKHSA publications number: GOV-10368



You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v3.0. To view this licence, visit [OGL](#). Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned.



UKHSA supports the UN
Sustainable Development Goals

