

Response to draft SACN Report on Carbohydrates and Health Report

Comments Related to Resistant Starch

August 2014

Effect of Resistant Starch on Post-Prandial Glycaemia

We welcome SACN recognition that resistant starch is considered as dietary fibre on the basis of its capacity to increase faecal mass. However, there is also adequate evidence for an effect of resistant starch on reducing post-prandial glycaemia which is both beneficial with higher proportions of resistant starch and biologically relevant. It is noted that only fasting blood glucose concentrations have been evaluated as a health benefit in the report and post-prandial values, which are listed by SACN as a physiological benefit, have been excluded. We are therefore of the opinion that the report should include an effect of resistant starch on post-prandial glycaemia. Please find below comments and relevant data references to support and substantiate this position:-

1. **Sections 9.35 to 9.49** discuss the effect of resistant starch on the following outcomes: energy intake, faecal weight, faecal pH and short-chain fatty acid content and faecal bacteria. The evidence for an effect of resistant starch on post-prandial glycaemia was not reviewed, despite there being numerous studies which address this research question. This omission is surprising since this evidence was recently reviewed by EFSA who concluded that **“a cause and effect relationship has been established between the consumption of resistant starch from all sources, when replacing digestible starch in baked foods, and a reduction of post-prandial glycaemic responses.”**(EFSA NDA Panel 2011).

2. EFSA's conclusion was based on a review of 14 randomised controlled studies, 12 of which were published after 1990, the cut-off date used for the SACN Report (see Appendix A). The EFSA Panel noted that most of the studies reported a statistically significant decrease in post-prandial glycaemic and insulinaemic responses when resistant starch replaced digestible starch in test products such as muffins, bread and crackers. Based on the findings of a dose-response study, they concluded that significant effects on post-prandial glucoses responses were seen when 14% or more of total starch was replaced by resistant starch. This condition is certainly achievable through diets typically consumed in the UK with the use of appropriately substituted products.

3. The EFSA Panel considered that the reduction of post-prandial glycaemic responses may be a beneficial physiological effect with the proviso that post-prandial insulinaemic responses were not disproportionately increased. Further, SACN's previous position statement on Dietary Fibre included reducing post-prandial glycaemia as a beneficial physiological effect(SACN 2008) (Section 1.2, para 6). One randomised controlled trial of 20 insulin-resistant subjects has found that consumption of resistant starch for 12 weeks led to an improvement in insulin sensitivity(Johnston et al. 2010).

4. Following the grading system for judging the evidence set out in the draft SACN report, the evidence from 12 randomised controlled studies shows that there is adequate evidence of an effect of resistant starch on reducing post-prandial glycaemia which is both beneficial with higher proportions of resistant starch and biologically relevant. We can only surmise that this evidence was not considered by SACN because of the short timescale of these studies. However, post-prandial glycaemia is a physiological response to a meal and therefore studies to measure it must, by definition, be short term studies.

References

- Behall, K. M. and J. Hallfrisch (2002). "Plasma glucose and insulin reduction after consumption of breads varying in amylose content." *Eur J Clin Nutr* **56**(9): 913-20.
- Behall, K. M. and J. C. Howe (1995). "Effect of long-term consumption of amylose vs amylopectin starch on metabolic variables in human subjects." *Am J Clin Nutr* **61**(2): 334-340.
- Behall, K. M. and D. J. Scholfield (2005). "Food amylose content affects postprandial glucose and insulin responses." *Cereal Chemistry* **82**(6): 654-659.
- Behall, K. M., D. J. Scholfield and J. G. Hallfrisch (2006). "Barley b-glucan reduces plasma glucose and insulin responses compared with resistant starch in men." *Nutrition Research* **26**: 644-650.
- Behall, K. M., D. J. Scholfield, J. G. Hallfrisch and H. G. Liljeberg-Elmstahl (2006). "Consumption of both resistant starch and beta-glucan improves postprandial plasma glucose and insulin in women." *Diabetes Care* **29**(5): 976-81.
- Brighenti, F., L. Benini, D. Del Rio, C. Casiraghi, et al. (2006). "Colonic fermentation of indigestible carbohydrates contributes to the second-meal effect." *Am J Clin Nutr* **83**(4): 817-22.
- EFSA NDA Panel (2011). "Scientific Opinion on the substantiation of health claims related to resistant starch and reduction of post-prandial glycaemic responses (ID 681), "digestive health benefits" (ID 682) and "favours a normal colon metabolism" (ID 783) pursuant to Article 13(1) of Regulation (EC) No 1924/2006." *EFSA Journal* **9**(4): 2024.
- Higgins, J. A., D. R. Higbee, W. T. Donahoo, I. L. Brown, et al. (2004). "Resistant starch consumption promotes lipid oxidation." *Nutr Metab (Lond)* **1**(1): 8.
- Hoebler, C., A. Karinthe, H. Chiron, M. Champ, et al. (1999). "Bioavailability of starch in bread rich in amylose: Metabolic responses in healthy subjects and starch structure." *Eur J Clin Nutr* **53**(5): 360-366.
- Jenkins, D. J., V. Vuksan, C. W. Kendall, P. Wursch, et al. (1998). "Physiological effects of resistant starches on fecal bulk, short chain fatty acids, blood lipids and glycemic index." *J Am Coll Nutr* **17**(6): 609-16.
- Johnston, K. L., E. L. Thomas, J. D. Bell, G. S. Frost, et al. (2010). "Resistant starch improves insulin sensitivity in metabolic syndrome." *Diabet Med* **27**(4): 391-7.
- Muir, J., Z. Lu, G. Collier and K. O'Dea (1994). The Acute Effects of High Resistant Starch Bread (Made from Hi-maize - A High Amylose Maize Starch) on Glucose and Insulin Responses in Non-diabetics. *Resistant starch: Plant breedings, applications development and commercial use*. I. L. Brown, K. J. McNaught, D. Andres and T. Morita. Oxford, Engl, Blackwell Science Ltd: 401-412.
- Noakes, M., P. M. Clifton, P. J. Nestel, R. Le Leu, et al. (1996). "Effect of high-amylose starch and oat bran on metabolic variables and bowel function in subjects with hypertriglyceridemia." *Am J Clin Nutr* **64**(6): 944-951.
- SACN (2008). Statement on Dietary Fibre.
http://www.sacn.gov.uk/pdfs/final_sacn_position_statement_for_website_dietary_fibre.pdf.
- Weickert, M. O., M. Mohlig, C. Koebnick, J. J. Holst, et al. (2005). "Impact of cereal fibre on glucose-regulating factors." *Diabetologia* **48**(11): 2343-53.

Appendix A: References to support effect of Resistant Starch on Post-Prandial Glycaemia

Reference	Subjects (Gender; Age)	Number	Design	Length of inter- vention	Food vehicle (Control)	Amount RS2	Effect on post- prandial glucose levels	Effect on glucose AUC	Effect on post- prandial insulin levels	Effect on insulin levels
Muir et al, 1994	Healthy men and women 31 +/- 7 years	3 men; 8 women	crossover	2 hours	Bread (control - 0% RS bread)	2g/100g/RS	=	↑NS	↓NS	↓NS
						3.2g/100gRS	↓NS	↓NS	↓NS	↓NS
						5.6g/100gRS	↓p<0.05	↓p<0.05	↓NS	↓p<0.01
						6.6g/100g RS	NR	↓NS	NR	↓p<0.01
Behall and Howe 1995	Hyperinsulin- aemic(HI) men 28 – 58 years	24 men, 14 hyperinsulin aemic	crossover	14 weeks (test meal 3 hours)	Muffins, cookies, puddings	30% amylose vs 70% amylose (HI)	↓p<0.05	↓NS	↓p<0.01	↓p<0.05
						30% amylose vs 70% amylose (C)	↓NS	↓NS	↓p<0.01	↓NS
Noakes et al, 1996	Hypertriglycerid aemic 44 – 64 years	13 men; 10 women	crossover	4 weeks (test meal 1.75 hours)	Muffins (control – muffins 1.3g RS)	5.8g/RS	↓p<0.03	↓NS	↓p<0.001	↓p<0.01
Jenkins et al, 1998	Healthy men and women 22 – 53 years	12 men; 12 women	crossover	2 weeks (test meal 2 hours)	Muffins, cereal (control – muffins 2.3g RS)	21.5g RS	NR	=	NR	NR
Hoebler et al, 1999	Healthy men and women 21 – 28 years	6 men; 2 women	crossover	4 hours	Bread, pasta	16.5g RS	↓p<0.05	↓p<0.05	↓p<0.05	↓p<0.05
Behall and Hallfrisch 2002	Healthy men and women 23 – 58 years	13 men, 12 women	latin square	3 hours	Bread (control – glucose solution)	2.0g RS	↓p<0.03	↓p<0.001	=	↓NS
						3.8g RS	↓p<0.03	↓p<0.001	=	↓NS
						8.2g RS	↓p<0.03	↓p<0.001	↓NS	↓NS
						11.5g RS	↓p<0.03	↓p<0.001	↓p<0.05	↓p<0.001
						13.4g RS	↓p<0.03	↓p<0.001	↓p<0.05	↓p<0.001
Higgins et al, 2004	Healthy men and women 33 +/- 5 years	7 men; 5 women	crossover	24 hours (test meal 6 hours)	Muffins, pasta, drink, biscuit C- 0% RS	2.7% RS	=	↑NS	=	↑NS
						5.4% RS	=	↑NS	=	=
						10.7% RS	=	↑NS	=	↑NS

Behall and Scholfield 2005	Healthy men and women 25 – 57 years	12 men; 12 women	latin square	3 hours	Cornchips (Control – cornchips 0g RS)	11.9g RS	↓ _{p<0.001}	↓ _{p<0.05}	↓ _{p<0.001}	↓ _{p<0.05}
					Muffins (Control – muffins 0g RS)	24.7g RS	↓ _{p<0.001}	↓ _{p<0.05}	↓ _{p<0.001}	↓ _{p<0.05}
Weickert et al, 2005	Healthy women 23 +/- 2 years	9	crossover	5 hours	Bread (Control – white bread)	10.4g RS	=	↓ _{NS}	↑ _{NS}	↓ _{NS}
Behall et al, 2006	Overweight women 43 years	10 normal weight; 10 overweight	latin square	4 hours	Muffins (Control – muffin 0.9g RS)	3.4g RS	↓ _{p<0.05}	↓ _{NS}	↓ _{NS}	↓ _{NS}
						6.5g RS	↓ _{p<0.05}	↓ _{NS}	↓ _{NS}	↓ _{p<0.05}
Behall et al, 2006	Overweight men 25 – 56 years	10 normal weight; 10 overweight	latin square	4 hours	Muffins (Control – muffin 0.1gRS)	6.3g RS	↓ _{NS}	↓ _{NS}	↓ _{NS}	=
						12.7gRS	↓ _{NS}	↓ _{NS}	↓ _{NS}	=
Brighenti et al, 2006	Healthy men and women 40 +/- 10 years	8 men; 2 women	crossover	9 hours	Cake (Control – cake 1g RS)	13g RS	↓ _{p<0.05}	NR	↓ _{p<0.02}	NR
<u>Number of significant reductions + Number of non-significant reductions</u> <u>Total number of comparisons</u>							(14 + 4)/23	(9 + 10)/24	(9 + 8)/23	(10 + 8)/23

↓_{p<0.05}: significant reduction in blood glucose level at at least one time-point; ↓_{NS}: non-significant reduction in blood glucose levels; =: no difference in blood glucose levels; NR: not reported.; AUC: area under curve