

**Additional analyses performed subsequent to the original  
systematic reviews to inform the Carbohydrates Working  
Group deliberations.**

---

# Contents


<b><i>Additional analyses performed subsequent to the original systematic reviews to inform the Carbohydrates Working Group deliberations. ....</i></b>	<b><i>1</i></b>
<b><i>Additional colo-rectal health analyses .....</i></b>	<b><i>3</i></b>
Dietary fibre intake and the risk of colo-rectal cancer (report paragraphs 8.26-8.28).....	3
Dietary fibre intake and the risk of colon cancer (report paragraphs 8.29-8.30) .....	4
Dietary fibre intake and the risk of rectal cancer (report paragraphs 8.31-8.32) .....	5
Cereal fibre intake and the risk of colo-rectal cancer (report paragraphs 8.114-8.115) .....	6
Dietary fibre and colo-rectal cancer risk: adjusting studies that determine dietary fibre as NSP to approximate AOAC values (report paragraph A2.11) .....	7
Non-digestible oligosaccharide or inulin supplementation and calcium absorption in adults (report paragraphs 9.35-9.37).....	9
Non-digestible oligosaccharide or inulin supplementation and calcium absorption in children and adolescents (report paragraphs 9.68-9.69) .....	10
Investigation of the dose-response relationship between dietary fibre and faecal wet weight and, where possible, intestinal transit time .....	11
Non-digestible oligosaccharides (report paragraph 9.24) .....	11
Resistant starch (report paragraph 9.45) .....	13
Wheat fibre (report paragraph 8.99) .....	15
<b><i>Additional cardio-metabolic health analyses.....</i></b>	<b><i>21</i></b>
Meta-regression of trials investigating sugars intake in relation to energy intake (report paragraph A9.1) .....	21
Outcome data (weighted mean difference in total energy intake) .....	21
Exposure data (mean difference in percentage energy as sugars) .....	21
Difference in sugar intake calculations for each trial .....	22
Meta-regression analysis .....	24
Higher carbohydrate, lower fat diets and fasting total cholesterol:HDL-cholesterol ratio (report paragraphs 5.40-5.41) .....	25
Cereal fibre (excluding oat fibre) in relation to energy intake (report paragraphs 8.95-8.96) ....	26
Glycaemic index and glycaemic load trials (report paragraphs 10.6 – 10.54) .....	27
Glycaemic index trials .....	28
Glycaemic load trials .....	38
<b><i>References.....</i></b>	<b><i>46</i></b>

## Additional colo-rectal health analyses

### *Dietary fibre intake and the risk of colo-rectal cancer (report paragraphs 8.26-8.28)*

1. Further meta-analyses were performed which included more recent data from the EPIC cohort (Murphy et al 2012), as identified in the update search, and excluding the previous publication from the EPIC cohort that was included in the colo-rectal health review (Bingham et al 2005). The cancer results presented below have been scaled down in the Carbohydrates and Health report to an increment of 7g/day of dietary fibre.

**Figure 1. Forest plot of per unit analysis (10g/day) for dietary fibre intake and colo-rectal cancer risk**



**Table 1. Results of per unit meta-analysis (10g/day) for dietary fibre intake and colo-rectal cancer risk**

Model	Pooled risk ratio estimate <sup>1</sup>		
	No. <sup>2</sup>	RR (95%CI)	Z (p-value)
Random effect	11	0.89 (0.82-0.96)	-3.06 (p=0.002)


<sup>1</sup>  $I^2 = 48.0\%$ ; p for test of heterogeneity = 0.037

<sup>2</sup> No. of relative risk estimates included in the pooled analysis.

2. Scaled down the per-unit meta-analysis for 7g/d dietary fibre increase in relation to colo-rectal cancer risk: RR 0.92, 95% CI 0.87, 0.97; p=0.002.

## ***Dietary fibre intake and the risk of colon cancer (report paragraphs 8.29-8.30)***

**Figure 2. Forest plot of per unit analysis (10g/day) for dietary fibre intake and colon cancer risk**



**Table 2. Results of per unit meta-analysis (10g/day) for dietary fibre intake and colon cancer risk**

Model	Pooled risk ratio estimate <sup>1</sup>		
	No. <sup>2</sup>	RR (95%CI)	Z (p-value)
Random effect	9	0.90 (0.84-0.97)	-2.70 (p=0.007)

<sup>1</sup>  $I^2 = 34.2\%$ ; p for test of heterogeneity = 0.144

<sup>2</sup> No. of relative risk estimates included in the pooled analysis.

3. Scaled down the per-unit meta-analysis for 7g/d dietary fibre increase in relation to colon cancer risk: RR 0.93, 95% CI 0.89, 0.98; p=0.007.

## Dietary fibre intake and the risk of rectal cancer (report paragraphs 8.31-8.32)

Figure 3. Forest plot of per unit analysis (10g/day) for dietary fibre intake and rectal cancer risk




Table 3. Results of per unit meta-analysis (10g/day) for dietary fibre intake and rectal cancer risk

Model	Pooled risk ratio estimate <sup>1</sup>		
	No. <sup>2</sup>	RR (95%CI)	Z (p-value)
Random effect	8	0.88 (0.81-0.96)	-2.69 (p=0.007)


<sup>1</sup>  $I^2 = 12.1\%$ ; p for test of heterogeneity = 0.336

<sup>2</sup> No. of relative risk estimates included in the pooled analysis.

4. Scaled down the per-unit meta-analysis for 7g/d dietary fibre increase in relation to rectal cancer risk: RR 0.91, 95% CI 0.86, 0.97; p=0.007.

## Cereal fibre intake and the risk of colo-rectal cancer (report paragraphs 8.114-8.115)

**Figure 4. Forest plot of per unit analysis (10g/day) for cereal fibre intake and colo-rectal cancer risk**



**Table 4. Results of per unit meta-analysis (10g/day) for cereal fibre intake and colo-rectal cancer risk**

Model	Pooled risk ratio estimate <sup>1</sup>		
	No. <sup>2</sup>	RR (95%CI)	Z (p-value)
Random effect	8	0.89 (0.84-0.94)	-3.12 (p<0.001)

<sup>1</sup>  $I^2 = 0.00\%$ ; p for test of heterogeneity = 0.666


<sup>2</sup> No. of relative risk estimates included in the pooled analysis.

5. Scaled down the per-unit meta-analysis for 7g/d cereal fibre increase in relation to colo-rectal cancer risk: RR 0.92, 95% CI 0.89, 0.96; p=0.001.

***Dietary fibre and colo-rectal cancer risk: adjusting studies that determine dietary fibre as NSP to approximate AOAC values (report paragraph A2.11)***

6. One study stated that dietary fibre was determined as NSP (Pietinen *et al.*, 1999). This provided an estimate for total colo-rectal cancer risk in relation to dietary fibre intakes. This study did not report colon and rectal cancer separately. The RR from this study was scaled down by a factor of 1.3 to give 10 g AOAC value from a 10 g NSP value.
7. In the EPIC cohort study (Bingham *et al.*, 2005; Murphy *et al.*, 2012) the AOAC method was used for all countries, except in the UK and Greece, where the NSP method was used. The fibre variable used in their analyses was obtained from the EPIC nutrient data base; in which the nutritional composition of foods across the different countries had been standardised. The earlier EPIC report (Bingham *et al.*, 2005) was used for analyses of cereal, vegetable, fruit and legume fibre in relation to colo-rectal cancer, as the later paper (Murphy *et al.*, 2012) did not report on these dietary fibre constituents. It is not possible to adjust the dietary fibre intake data within the EPIC cohort study.
8. Only the meta-analysis of dietary fibre intake in relation to total colo-rectal cancer risk needed to be re-analysed to adjust NSP for AOAC values for Pietinen *et al.*, 1999. Figure 5 and Table 5 **Error! Reference source not found.** show the forest plot and meta-analysis with the adjusted intake data. The assigned weight to the Pietinen *et al.*, 1999 study increases slightly in the adjusted analysis, as the variation is reduced due to the RR being scaled down by a factor of 1.3. This has very little impact on the overall pooled risk ratio estimate (compare with Figure 1 and Table 1 above).

**Figure 5 Forest plot of per unit analysis (10g AOAC/day) for dietary fibre intake and colo-rectal cancer risk**



**Table 5. Results of per unit meta-analysis (10g AOAC/day) for dietary fibre intake and colo-rectal cancer risk**

Model	Pooled risk ratio estimate <sup>1</sup>		
	No. <sup>2</sup>	RR (95%CI)	Z (p-value)
Random effect	11	0.89 (0.83-0.96)	-2.89 (p=0.003)

<sup>1</sup>  $I^2 = 54.9\%$ ; p for test of heterogeneity = 0.014


<sup>2</sup> No. of RR estimates included in pooled analysis.

9. Scaled down the per-unit meta-analysis for 7g/d dietary fibre increase in relation to colo-rectal cancer risk: RR 0.92, 95% CI 0.88, 0.97; p=0.003. This has very little impact on the overall pooled risk ratio estimate (compare with Figure 1 and Table 1 above).

## ***Non-digestible oligosaccharide or inulin supplementation and calcium absorption in adults (report paragraphs 9.35-9.37)***

10. Meta-analyses conducted based on the evidence detailed in Chapter 9.

**Figure 6. Forest plot of the effect of non-digestible oligosaccharides or inulin supplementation on fractional calcium absorption in adults**



**Table 6. Results of meta-analysis for non-digestible oligosaccharides or inulin supplementation on calcium absorption in adults**


Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	% (95%CI)	Z (p-value)
Random effect	4	0.47 (-3.36-4.29)	0.24(p=0.81)

<sup>1</sup>  $I^2 = 0.00\%$  (95% CI 0.00-84.69%); p for test of heterogeneity = 0.489

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

## Non-digestible oligosaccharide or inulin supplementation and calcium absorption in children and adolescents (report paragraphs 9.68-9.69)

**Figure 7. Forest plot of the effect of non-digestible oligosaccharides or inulin supplementation on fractional calcium absorption in children and adolescents**



**Table 7. Results of meta-analysis for non-digestible oligosaccharides or inulin supplementation on calcium absorption in children and adolescents**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	% (95%CI)	Z (p-value)
Random effect	5	4.95 (1.62-8.27)	2.726(p=0.003)


<sup>1</sup>  $I^2 = 51.79\%$  (95% CI 0.00-82.29%); p for test of heterogeneity = 0.081

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

# Investigation of the dose-response relationship between dietary fibre and faecal wet weight and, where possible, intestinal transit time


## Non-digestible oligosaccharides (report paragraph 9.24)

Figure 8. Sub-group analysis of non-digestible oligosaccharide (FOS, GOS & inulin) intake and faecal weight by exposure category



11. The test for heterogeneity between sub-groups is not significant ( $p=0.115$ ), indicating no significant dose-response relationship in the data.

**Figure 9. Meta-regression plot and analysis of non-digestible oligosaccharide (FOS, GOS & inulin) intake in relation to faecal wet weight**



Meta-regression  
REML estimate of between-study variance  
% residual variation due to heterogeneity  
Proportion of between-study variance explained  
with Knapp-Hartung modification


Number of obs = 11  
tau2 = 25.87  
I-squared\_res = 0.00%  
Adj R-squared = -44.67%

md	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
NDO	1.81137	1.150715	1.57	0.150	-.7917282	4.414469
_cons	-9.918389	15.85427	-0.63	0.547	-45.78324	25.94646

12. The meta-regression analysis is not significant ( $p=0.150$ ), indicating no significant linear dose-response relationship in the data. A limitation with the meta-regression analysis is that it fits a linear regression line, but does not determine non-linear relationship


## Resistant starch (report paragraph 9.45)

**Figure 10. Sub-group analysis of resistant starch (RS1, 2 & 3) intake and faecal weight by exposure category**



13. The test for heterogeneity between subgroups is not significant ( $p=0.689$ ), indicating no significant dose-response relationship in the data.

**Figure 11. Meta-regression plot and analysis of resistant starch (RS1, 2 & 3) intake in relation to faecal wet weight**



Meta-regression					Number of obs	=	12
REML estimate of between-study variance					tau2	=	0
% residual variation due to heterogeneity					I-squared_res	=	0.00%
Proportion of between-study variance explained					Adj R-squared	=	0.00%
With Knapp-Hartung modification							
md	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]		
RS	.2294103	.817321	0.28	0.785	-1.591694	2.050515	
_cons	33.88574	21.86565	1.55	0.152	-14.83397	82.60544	

14. The meta-regression analysis is not significant (p=0.785), indicating no significant linear dose-response relationship in the data.


## Wheat fibre (report paragraph 8.99)

15. In the trials measuring faecal wet weight several different analytical techniques were used to determine the dietary fibre content of wheat fibre. All dietary fibre values for wheat fibre have been converted to AOAC values based on published comparisons between the different techniques. The dietary fibre analytical techniques other than AOAC used in the trials were crude fibre, neutral detergent fibre (Van Soest), Southgate method and NSP. Values were converted to NSP then to AOAC values.
16. Neutral detergent fibre (Van Soest) determines insoluble fibre only and crude fibre determines part of the insoluble fibre. For kidney beans the crude fibre measure is 0.6 that of the neutral detergent fibre (Lunn & Buttriss, 2007). For the conversion of wheat fibre/bran as measured by crude fibre a ratio of 0.43 to NSP was used (AWT, 2005). The values from the McCance and Widdowson food composition tables 5<sup>th</sup> edition provide Southgate and NSP dietary fibre values for wheat bran and the 7<sup>th</sup> edition provides NSP and AOAC dietary fibre values for wheat bran. For wheat bran the Southgate value is 1.089 that for NSP and the NSP value is 0.799 that of the AOAC value. The values have been adjusted accordingly to estimate the AOAC value for wheat fibre in each trial, where this is not given.

AWT, Arbeitsgemeinschaft Wirkstoffe in der Tierernährung (2005) *Enzyme in der Tierernährung* (Enzymes in Animal Nutrition), 55pp.


Lunn, J & Buttriss, JL (2007) Carbohydrates and dietary fibre. *British Nutrition Foundation Nutrition Bulletin*, **32**, 21–64

**Figure 12. Sub-group analysis of wheat fibre intake and faecal weight by exposure category (AOAC values)**



17. The forest plot shows the individual trials in order of ascending dose (right column) and grouped into subgroups of <10g/day, 10-15g/day, >15g-20g/day and >20g/day AOAC dietary fibre. The dose is the mean difference in intake between intervention and control group.
18. The test for heterogeneity between sub-groups is significant ( $p < 0.001$ ), indicating a significant dose-response relationship in the data. This is investigated further in the meta-regression analysis below.


Figure 13. Meta-regression plot and analysis of wheat fibre intake in relation to faecal wet weight



Meta-regression					Number of obs	=	23
REML estimate of between-study variance					tau2	=	489.7
% residual variation due to heterogeneity					I-squared_res	=	61.85%
Proportion of between-study variance explained with Knapp-Hartung modification					Adj R-squared	=	69.71%
md	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]		
dose	4.84749	.8639021	5.61	0.000	3.050907	6.644073	
_cons	-12.5989	17.42783	-0.72	0.478	-48.84205	23.64424	


19. The meta-regression analysis is significant ( $p < 0.001$ ), indicating a significant linear dose-response relationship in the data. A 1g increase in wheat fibre intake results in a 4.8g (95%CI 3.0, 6.6) increase in faecal weight.
20. The colo-rectal health review estimated the effect on faecal wet weights as broadly equating to a 4g increase in faecal wet weight per 1g wheat fibre.

**Figure 14. Forest plot showing the effect of wheat fibre on intestinal transit time, in order of descending mean control intestinal transit time**




21. The forest plot shows the individual trials in order of descending mean control intestinal transit time (right column). The effect of wheat fibre on decreasing intestinal transit time is significant ( $p < 0.001$ ), but the heterogeneity is high ( $I^2 = 67\%$ ) and this is due to variation in the initial transit time (as determined by control values) modifying the response to wheat fibre. This is investigated further in the meta-regression analysis below.

**Figure 15. Sub-group analysis of randomised controlled trials investigating the modifying effect of initial transit time on the effect of wheat fibre on intestinal transit time**



22. A sub-group analysis within the meta-analysis was performed by categorising the differences in intestinal transit time into four groups based on initial transit time (>75 hours; 55-75 hours; 48-55 hours; and 30-47 hours). The test for heterogeneity between sub-groups was significant ( $p < 0.001$ ), indicating a significant modifying effect of initial transit time on the effect of wheat fibre on intestinal transit time.

**Figure 16. Meta-regression showing the modifying effect of initial transit time on the effect of wheat fibre on intestinal transit time**



Meta-regression					Number of obs	=	15
REML estimate of between-study variance					tau2	=	0
% residual variation due to heterogeneity					I-squared_res	=	0.00%
Proportion of between-study variance explained					Adj R-squared	=	100.00%
Joint test for all covariates					Model F(2,12)	=	19.15
With Knapp-Hartung modification					Prob > F	=	0.0002

md	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
dose	-.0791916	.2139272	-0.37	0.718	-.5452988	.3869157
mean0	-.6184224	.0999872	-6.19	0.000	-.8362759	-.4005689
_cons	20.82544	7.22623	2.88	0.014	5.080841	36.57005

23. The meta-regression includes two variables: wheat fibre dose and control group values (variable mean0 above). It is only possible to graph one variable, which is shown above for mean control intestinal transit time in relation to the effect of wheat fibre on total intestinal transit time. There is no linear dose-response relationship between wheat fibre and transit time in the data ( $p=0.718$ ), but the effect of initial transit time (as determined by mean control group values) on the intestinal transit time response to wheat fibre is significant ( $p<0.001$ ).
24. There are insufficient trials to enable the dose-response relationship to be investigated within a sub-set of trials in which subjects have a similar initial intestinal transit time.

## **Additional cardio-metabolic health analyses**

### ***Meta-regression of trials investigating sugars intake in relation to energy intake (report paragraph A9.1)***

25. The exposure measure (mean difference in percentage energy intake as sugars) and outcome measure (mean difference in total energy intake) are inter-related. A meta-regression of eleven trials investigating sugars intake in relation to energy intake was performed using exposure data as a comparison of end of intervention if the outcome measure with variance data was reported thus and change from baseline if the outcome measure with variance data was reported thus.

### **Outcome data (weighted mean difference in total energy intake)**

26. The outcome measure with variance data were those reported in the trial, except for Drummond *et al.*, 2003, which was analysed using imputed variance data for change from baseline value comparisons (to avoid misrepresentation of results – see Annex 9 in report). Except for that one trial, this approach did not require imputation of further variance data.

### **Exposure data (mean difference in percentage energy as sugars)**

27. The mean difference in sugars intake was calculated from a comparison of the end of intervention data in seven trials (Drummond & Kirk, 1998; Poppitt *et al.*, 2002; Raben *et al.*, 2002; Brynes *et al.*, 2003; Reid *et al.*, 2010; Aeberli *et al.*, 2011; Reid *et al.*, 2014). Two trials only reported the amount of sugars subjects were supplemented with and this was assumed to represent the difference in intakes between the control and intervention groups (Reid *et al.*, 2007; Njike *et al.*, 2011). For two trials the mean difference in sugars intake was calculated from a comparison of change from baseline data (Saris *et al.*, 2000; Drummond *et al.*, 2003).
28. Four trials involved multiple interventions and only the most comparable groups, with the exception of sugars intake, have been used in the analyses (Poppitt *et al.*, 2002; Brynes *et al.*, 2003; Aeberli *et al.*, 2011; Njike *et al.*, 2011). In one trial the intervention groups comprised a high and low GI group, a high fat and high sucrose group (Brynes *et al.*, 2003). The data from the high sucrose group were compared to the high GI group, as these two groups had the most similar dietary intakes with the exception of sucrose intake – the low GI diet had a higher dietary fibre content and the high fat had a lower carbohydrate content. Another trial compared moderate fructose, moderate glucose, high fructose, high glucose, high sucrose and low fructose intervention groups (Aeberli *et al.*, 2011). For the analyses the outcome/exposure data from the high sucrose group were compared to the low fructose group, as these two groups had the greatest difference in sugars intake and there was no specific low sucrose group. The data for sucrose, free glucose and free fructose were combined to give sugars exposure values for each group.
29. A third trial comprised a control diet, a low-fat, complex carbohydrate diet and a low-fat, simple carbohydrate diet (Poppitt *et al.*, 2002). For the analyses the

outcome/exposure data from the low-fat, simple carbohydrate group were compared to the low-fat, complex carbohydrate group, as these two groups had the most similar dietary intakes with the exception of sugars intake. A fourth trial comprised a control diet, a sugar-free cocoa and a sugar-sweetened cocoa group (Njike *et al.*, 2011). For the analyses the outcome/exposure data from the sugar-free cocoa group were compared to the sugar-sweetened cocoa group, as these two groups had the most similar dietary intakes with the exception of sugars intake.

30. To convert grams of sugar to percentage energy. values have been used for individual sugars as follows: 15.7 kJ/g glucose, 15.2kJ/g fructose, 16.3 kJ/g sucrose (Elia & Cummings, 2007). In two trials, which used sucrose as the intervention, only the weight for total sugars intake was reported (Reid *et al.*, 2010; Reid *et al.*, 2014); a value of 16.0 kJ/g was used to convert grams of total sugar to percentage energy.

### **Difference in sugar intake calculations for each trial**

(Drummond & Kirk, 1998)

End of intervention data

% energy NMES 8.1 vs. 10

Difference = 1.9 % total energy

(Saris *et al.*, 2000)

Only change from baseline data reported

% energy simple carbohydrates 7.2 vs. -3.5

Difference = 10.7 % total energy

(Poppitt *et al.*, 2002)

End of intervention data

Six month combined intervention data used for outcome measure as this was the only measure with variance; therefore, six month combined data used for exposure measure.

% energy simple carbohydrates 17.6 vs. 28.9

Difference = 11.3 % total energy

(Raben *et al.*, 2002)

End of intervention data

% energy sucrose 27 vs. 4

Difference = 23 % total energy

(Brynes *et al.*, 2003)

End of intervention data

Sucrose 132g vs. 46g

$132 \times 16.3\text{kJ} = 2.15\text{MJ}$ ;  $46 \times 16.3\text{kJ} = 0.75\text{MJ}$

Difference =  $(2.15/9.9 \times 100) - (0.75/9.02 \times 100) = 13.4 \%$  total energy

(Drummond *et al.*, 2003)

Change from baseline data with variance computed

% energy NMES Change from 10 to 10.5 vs. Change from 11.4 to 9.0: 0.5 vs. -2.4

Difference = 2.9% total energy

(Reid *et al.*, 2007)

End of intervention data. No sugars intake data, but the sucrose supplements provided 105g/day sucrose  
 $105\text{g} \times 16.3\text{kJ/g} = 1.71\text{ MJ/day}$ .  
End of intervention total energy intake of sucrose group = 8.722MJ/day  
Assumed difference =  $1.71/8.722 \times 100 = 19.6\%$  total energy

(Reid *et al.*, 2010)

End of intervention data  
Total sugars 196.65g vs. 105.06g  
 $196.65 \times 16.0\text{kJ} = 3.15\text{MJ}$ ;  $105.06 \times 16.0\text{kJ} = 1.68\text{MJ}$   
Difference =  $(3.15/9.3 \times 100) - (1.68/8.22 \times 100) = 13.4\%$  total energy

(Aeberli *et al.*, 2011)

End of intervention data  
The data for sucrose, free glucose and free fructose were combined to give sugars intake  
Low fructose =  $(53.1\text{g} \times 16.3\text{kJ}) + (8\text{g} \times 15.7\text{kJ}) + (7.2\text{g} \times 15.2\text{kJ}) = 1.10\text{MJ}$   
High sucrose =  $(130.4\text{g} \times 16.3\text{kJ}) + (14\text{g} \times 15.7\text{kJ}) + (13.4 \times 15.2\text{kJ}) = 2.55\text{ MJ}$   
Difference =  $(2.55/10.86 \times 100) - (1.10/9.79 \times 100) = 12.2\%$  total energy

(Njike *et al.*, 2011)

End of intervention data. No sugars intake data, but the sucrose-sweetened drinks provided 91 g/d  
 $91\text{g} \times 16.3\text{kJ} = 1.48\text{MJ}$   
End of intervention total energy intake of sucrose group =  $1991.4\text{ kcal} \times 4.184 = 8.33\text{MJ/day}$   
Assumed difference =  $1.48/8.33 \times 100 = 17.8\%$  total energy


(Reid *et al.*, 2014)

End of intervention data  
Total sugars 177.5g vs. 88.4g  
 $177.5 \times 16.0\text{kJ} = 2.84\text{MJ}$ ;  $88.4 \times 16.0\text{kJ} = 1.41\text{MJ}$   
Difference =  $(2.84/9.091 \times 100) - (1.41/7.996 \times 100) = 13.55\%$  total energy

Meta-regression analysis

31. The regression coefficient obtained from a meta-regression analysis describes how the outcome variable (weighted mean difference in total energy intake; kJ/day) changes with a unit increase in the explanatory variable (mean difference in % energy from sugars/day).
32. The meta-regression analysis shows a significant linear dose-response relationship with a 69kJ (95%CI 3,135kJ; p=0.042) change in total energy intake per one percentage change in energy from sugars (see **Error! Not a valid bookmark self-reference.**).


Figure 17. Meta-regression plot and analysis of trials investigating sugars intake in relation to energy intake using exposure data as a comparison of end of intervention if the outcome measure with variance data was reported thus and change from baseline if the outcome measure with variance data was reported thus



### ***Higher carbohydrate, lower fat diets and fasting total cholesterol:HDL-cholesterol ratio (report paragraphs 5.40-5.41)***

33. The mean difference values used in the Cardio-metabolic review were incorrect for one of the trials (Howard *et al.*, 2006). In the review the value on the forest plot (figure 2.43) for the mean difference for the Howard et al 2006 paper is reported as -0.10 95% CI -0.15, -0.03, but in the paper it is reported as -0.04, 95% CI -0.13, 0.5. Therefore, the data have been re-analysed and the pooled estimate given below has been used in the Carbohydrates and Health report.

***Figure 18. Forest plot displaying randomised controlled trials investigating the effect of higher carbohydrate, lower fat diets on fasting total cholesterol:HDL-cholesterol ratio***



***Table 8. Results of meta-analysis for higher carbohydrate, lower fat diets on fasting total cholesterol:HDL-cholesterol ratio***

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD (95%CI)	Z (p-value)
Random effect	5	-0.03 (-0.12, 0.05)	0.75 (p=0.455)


<sup>1</sup> I<sup>2</sup> = 0.0%; p for test of heterogeneity = 0.734

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

## ***Cereal fibre (excluding oat fibre) in relation to energy intake (report paragraphs 8.95-8.96)***

34. One randomised controlled trial included in the original meta-analysis examined the relationship between cocoa bran and energy intake (Jenkins *et al.*, 2000). As cocoa bran is a legume fibre and not a cereal fibre the meta-analysis was performed with this trial excluded. Four randomised controlled trials were included in a meta-analysis (Tredger *et al.*, 1991; Sanders & Reddy, 1992; Jenkins *et al.*, 1999; Vuksan *et al.*, 1999).
35. No significant effect is demonstrated for cereal fibre (excluding oat fibre) consumption on energy intake (-269kJ, 95% CI (-826, 288 kJ; p=0.34).

**Figure 19. Forest plot displaying randomised controlled trials investigating cereal fibre intake (excluding oat fibre) in relation to energy intake (KJ/day)**



**Table 9. Results of meta-analysis for cereal fibre intake (excluding oat fibre) in relation to energy intake (KJ/day)**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD (95%CI)	Z (p-value)
Random effect	4	-269KJ (-826, 288)	0.95 (p=0.344)

<sup>1</sup> I<sup>2</sup> = 0.0%; p for test of heterogeneity = 0.423

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.


***Glycaemic index and glycaemic load trials (report paragraphs 10.6 – 10.54)***

36. In the cardio-metabolic health review, trials investigating glycaemic index (GI) and glycaemic load (GL) were combined into a single meta-analysis for each health outcome. To assist with interpreting the evidence, the trials were subsequently categorised into GI and GL studies and included in separate meta-analyses. Details of the conclusions drawn from these analyses can be found in Chapter 10 of the Carbohydrates and Health report.
37. The difference between these two types of trial is that the glycaemic index trials do not vary carbohydrate quantity, but change the quality to modify the GI. The GL trials reduce carbohydrate intake, resulting in an increased proportion of fat, including saturated fatty acids, and/or protein intake, as well as changing the carbohydrate quality to modify the glycaemic index. Both dietary strategies modify glycaemic index and glycaemic load, but the latter strategy modifies glycaemic load more and the former glycaemic index more.
38. In relation to the fasting lipid concentrations, one trial reported these as change from baseline (McMillan-Price *et al.*, 2006), whereas all other trials reported follow-up values. In the meta-analyses and corresponding forest plots, a positive number on the x-axis indicates that high GI/GL is greater than low GI/GL. For the weight loss meta-analyses and corresponding forest plots, both experimental groups have negative weight values, i.e. weight loss, and a positive value means less weight loss in the high GI/GL group.

## Glycaemic index trials

### Blood pressure (report paragraphs 10.12-10.13)

**Figure 20. Forest plot for glycaemic index diets and diastolic blood pressure (mmHg)**




**Table 10. Results of meta-analysis for low glycaemic index diet vs. high glycaemic index diet and diastolic blood pressure.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmHg (95%CI)	Z (p-value)
Random effect	4	0.60 (-2.06-3.25)	0.44 (p=0.660)

<sup>1</sup>  $I^2 = 22.8\%$ ; p for test of heterogeneity = 0.274

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

**Figure 21. Forest plot for glycaemic index diets and systolic blood pressure (mmHg)**



**Table 11. Results of meta-analysis for low glycaemic index diet vs. high glycaemic index diet and systolic blood pressure.**


Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmHg (95%CI)	Z (p-value)
Random effect	4	-0.54 (-4.08-2.99)	0.30 (p=0.764)

<sup>1</sup>  $I^2 = 28.4\%$ ; p for test of heterogeneity = 0.242

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

## Fasting blood lipids (report paragraphs 10.14-10.20)

**Figure 22. Forest plot for glycaemic index diets and fasting total cholesterol (mmol/L)**




**Table 12. Results of meta-analysis for low glycaemic index diet vs. high glycaemic index diet and fasting total cholesterol concentration.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	9	0.20 (0.08-0.33)	3.17 (p=0.002)

<sup>1</sup> I<sup>2</sup> = 23.4%; p for test of heterogeneity = 0.235

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

**Figure 23. Forest plot for glycaemic index diets and fasting LDL-cholesterol (mmol/L)**




**Table 13. Results of meta-analysis for low glycaemic index diet vs. high glycaemic index diet and fasting LDL-cholesterol concentration.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	9	0.21 (0.10-0.32)	3.71 (p=0.000)

<sup>1</sup>  $I^2 = 0\%$ ; p for test of heterogeneity = 0.708

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

**Figure 24. Forest plot of the total cholesterol and LDL-cholesterol trials above, showing mean difference in weight loss between low and high glycaemic index diets (kg) (report paragraph 10.16)**



**Table 14. Results of meta-analysis for low glycaemic index diet vs. high glycaemic index diet and mean difference in weight loss.**


Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	kg (95%CI)	Z (p-value)
Random effect	8	0.68 (-0.05-1.42)	1.81 (p=0.069)
Fixed effect	8	0.62 (0.10-1.13)	2.34 (p=0.019)

<sup>1</sup>  $I^2 = 47.60\%$ ; p for test of heterogeneity = 0.064

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

39. One trial does not report the weight loss from baseline, but gives total weights from which it is not possible to calculate the mean difference with variation. The difference in the mean weight loss values in this trial is -4.0kg for the low GI diet and -4.5kg for the high GI diet (Bellisle *et al.*, 2007).

**Figure 25. Forest plot for glycaemic index diets and fasting HDL-cholesterol (mmol/L)**




**Table 15. Results of meta-analysis for low glycaemic index diet vs. high glycaemic index diet and fasting HDL-cholesterol concentration.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	10	0.01 (-0.05-0.06)	0.24 (p=0.810)

<sup>1</sup> I<sup>2</sup> = 44.0%; p for test of heterogeneity = 0.066

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

**Figure 26. Forest plot for glycaemic index diets and fasting triacylglycerol (mmol/L)**




**Table 16. Results of meta-analysis for low glycaemic index diet vs. high glycaemic index diet and fasting triacylglycerol concentration.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	10	-0.04 (-0.17-0.10)	0.55 (p=0.586)

<sup>1</sup>  $I^2 = 59\%$ ; p for test of heterogeneity = 0.009

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

**Figure 27. Forest plot for glycaemic index diets and fasting total cholesterol:HDL-cholesterol ratio**




**Table 17. Results of meta-analysis for low glycaemic index diet vs. high glycaemic index diet and fasting total cholesterol:HDL-cholesterol ratio .**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD (95%CI)	Z (p-value)
Random effect	5	0.01 (-0.20-0.21)	0.07 (p=0.945)

<sup>1</sup>  $I^2 = 44.7\%$  (95% CI 0.00-72.78); p for test of heterogeneity = 0.124

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

**Figure 28. Forest plot for glycaemic index diets and fasting non-esterified fatty acids (mmol/L)**



**Table 18. Results of meta-analysis for low glycaemic index diet vs. high glycaemic index diet and fasting non-esterified fatty acid concentration.**


Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	4	0.01 (-0.05-0.08)	0.34 (p=0.736)

<sup>1</sup>  $I^2 = 30.7\%$  (95% CI 0.00-72.78); p for test of heterogeneity = 0.228

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

## Metabolic measures (report paragraphs 10.27-10.28)

**Figure 29. Forest plot for glycaemic index diets and fasting blood glucose concentration (mmol/L)**



**Table 19. Results of meta-analysis for low glycaemic index diet vs. high glycaemic index diet and fasting blood glucose concentration.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	11	-0.01 (-0.09-0.07)	0.20 (p=0.845)


<sup>1</sup> I<sup>2</sup> = 1.9%; p for test of heterogeneity = 0.423

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

## Glycaemic load trials

### Blood pressure (report paragraphs 10.37-10.38)

**Figure 30. Forest plot for glycaemic load diets and diastolic blood pressure (mmHg)**




**Table 20. Results of meta-analysis for low glycaemic load diet vs. high glycaemic load diet and diastolic blood pressure.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmHg (95%CI)	Z (p-value)
Random effect	3	3.10 (0.25-5.95)	2.13 (p=0.033)

<sup>1</sup>  $I^2 = 0.0\%$ ; p for test of heterogeneity = 0.839

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

**Figure 31. Forest plot for glycaemic load diets and systolic blood pressure (mmHg)**



**Table 21. Results of meta-analysis for low glycaemic load diet vs. high glycaemic load diet and systolic blood pressure.**


Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmHg (95%CI)	Z (p-value)
Random effect	3	1.92 (-0.76-4.60)	1.40 (p=0.161)

<sup>1</sup> I<sup>2</sup> = 0.0%; p for test of heterogeneity = 0.455

<sup>2</sup> No. of MD estimates included in pooled analysis.

## Fasting Blood lipids (report paragraphs 10.39-10.40)

**Figure 32. Forest plot for glycaemic load diets and fasting total cholesterol (mmol/L)**




**Table 22. Results of meta-analysis for low glycaemic load diet vs. high glycaemic load diet and fasting total cholesterol concentration.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	3	-0.01 (-0.14-0.13)	0.08 (p=0.936)

<sup>1</sup> I<sup>2</sup> = 0%; p for test of heterogeneity = 0.862

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

**Figure 33. Forest plot for glycaemic load diets and fasting LDL-cholesterol (mmol/L)**




**Table 23. Results of meta-analysis for low glycaemic load diet vs. high glycaemic load diet and fasting LDL-cholesterol concentration.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	5	0.07 (-0.05-0.19)	1.06 (p=0.287)

<sup>1</sup>  $I^2 = 0\%$ ; p for test of heterogeneity = 0.416

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

**Figure 34. Forest plot for glycaemic load diets and fasting HDL-cholesterol (mmol/L)**




**Table 24. Results of meta-analysis for low glycaemic load diet vs. high glycaemic load diet and fasting HDL-cholesterol concentration.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	5	-0.01 (-0.18-0.16)	0.06 (p=0.951)

<sup>1</sup> I<sup>2</sup> = 82.9%; p for test of heterogeneity = 0.000

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

**Figure 35. Forest plot for glycaemic load diets and fasting triacylglycerol (mmol/L)**



**Table 25. Results of meta-analysis for low glycaemic load diet vs. high glycaemic load diet and fasting triacylglycerol concentration.**


Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	5	0.13 (0.03-0.24)	2.53 (p=0.012)

<sup>1</sup> I<sup>2</sup> = 33.1%; p for test of heterogeneity = 0.201

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

One trial reported on glycaemic load diets and fasting total cholesterol:HDL-cholesterol ratio (Maki *et al.*, 2007) and no trials reported on fasting non-esterified fatty acids.

**Figure 36. Forest plot of trials above, showing mean difference in weight loss between low and high glycaemic load diets (kg)**



**Table 26. Results of meta-analysis for low glycaemic load diet vs. high glycaemic load diet and mean difference in weight loss.**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	kg (95%CI)	Z (p-value)
Random effect	5	0.32 (-0.43-1.06)	0.82 (p=0.41)


<sup>1</sup>  $I^2 = 0.0\%$ ; p for test of heterogeneity = 0.655

<sup>2</sup> No. of mean difference estimates included in the pooled analysis.

N.B. data extracted from a figure for one trial (Ebbeling *et al.*, 2007)

## Metabolic measures (report paragraphs 10.47-10.48)

**Figure 37. Forest plot for glycaemic load diets and fasting blood glucose concentration (mmol/L)**



**Table 27. Results of meta-analysis for low glycaemic load diet vs. high glycaemic load diet and fasting blood glucose concentration).**

Model	Pooled mean difference estimate <sup>1</sup>		
	No. <sup>2</sup>	MD mmol/l (95% CI)	Z (p-value)
Random effect	3	-0.13 (-0.40-0.13)	1.40 (p=0.328)

<sup>1</sup>  $I^2 = 73.9\%$ ; p for test of heterogeneity = 0.022

<sup>2</sup> No. of MD estimates included in pooled analysis.

## References

- Abete I, Parra D & Martinez JA (2008) Energy-restricted diets based on a distinct food selection affecting the glycemic index induce different weight loss and oxidative response. *Clinical Nutrition* **27**, 545-551.
- Abrams SA, Griffin IJ, Hawthorne KM, Liang L, Gunn SK, Darlington G & Ellis KJ (2005) A combination of prebiotic short- and long-chain inulin-type fructans enhances calcium absorption and bone mineralization in young adolescents. *Am J Clin Nutr* **82**, 471-476.
- Aeberli I, Gerber PA, Hochuli M, Kohler S, Haile SR, Gouni-Berthold I, Berthold HK, Spinass GA & Berneis K (2011) Low to moderate sugar-sweetened beverage consumption impairs glucose and lipid metabolism and promotes inflammation in healthy young men: a randomized controlled trial. *American Journal of Clinical Nutrition* **94**, 479-485.
- Bellisle F, Dalix AM, De Assis MA, Kupek E, Gerwig U, Slama G & Oppert JM (2007) Motivational effects of 12-week moderately restrictive diets with or without special attention to the Glycaemic Index of foods. *British Journal of Nutrition* **97**, 790-798.
- Brynes AE, Mark EC, Ghatei MA, Dornhorst A, Morgan LM, Bloom SR & Frost GS (2003) A randomised four-intervention crossover study investigating the effect of carbohydrates on daytime profiles of insulin, glucose, non-esterified fatty acids and triacylglycerols in middle-aged men. *British Journal of Nutrition* **89**, 207-218.
- Colette C, Percheron C, Pares-Herbute N, Michel F, Pham TC, Brillant L, Descomps B & Monnier L (2003) Exchanging carbohydrates for monounsaturated fats in energy-restricted diets: effects on metabolic profile and other cardiovascular risk factors. *International Journal of Obesity* **27**, 648-656.
- Das SK, Gilhooly CH, Golden JK, *et al.* (2007) Long-term effects of 2 energy-restricted diets differing in glycemic load on dietary adherence, body composition, and metabolism in CALERIE: a 1-y randomized controlled trial. *American Journal of Clinical Nutrition* **85**, 1023-1030.
- Drummond S & Kirk T (1998) The effect of different types of dietary advice on body composition in a group of Scottish men. *Journal of Human Nutrition and Dietetics* **11**, 473-485.
- Drummond S, Kirk T, Jackson J, Hendry J, Panton S & Gray F (2003) Effectiveness of dietary advice given by community dietitians to men with elevated blood cholesterol in a clinical setting: a pilot study. *Journal of Human Nutrition and Dietetics* **16**, 81-83.
- Ebbeling CB, Leidig MM, Sinclair KB, Seger-Shippie LG, Feldman HA & Ludwig DS (2005) Effects of an ad libitum low-glycemic load diet on cardiovascular disease risk factors in obese young adults. *American Journal of Clinical Nutrition* **81**, 976-982.
- Ebbeling CB, Leidig MM, Feldman HA, Lovesky MM & Ludwig DS (2007) Effects of a

low-glycemic load vs low-fat diet in obese young adults: a randomized trial.[erratum appears in JAMA. 2007 Aug 8;298(6):627]. *JAMA* **297**, 2092-2102.

Elia M & Cummings JH (2007) Physiological aspects of energy metabolism and gastrointestinal effects of carbohydrates. *Eur J Clin Nutr* **61**, S40-S74.

Ginsberg HN, Kris-Etherton P, Dennis B, *et al.* (1998) Effects of reducing dietary saturated fatty acids on plasma lipids and lipoproteins in healthy subjects - The Delta Study, protocol 1. *Arterioscler Thromb Vasc Biol* **18**, 441-449.

Griffin IJ, Davila PM & Abrams SA (2002) Non-digestible oligosaccharides and calcium absorption in girls with adequate calcium intakes. *British Journal of Nutrition* **87 Suppl 2**, S187-S191.

Griffin IJ, Hicks PMD, Heaney RP & Abrams SA (2003) Enriched chicory inulin increases calcium absorption mainly in girls with lower calcium absorption. *Nutrition Research* **23**, 901-909.

Holloway L, Moynihan S, Abrams SA, Kent K, Hsu AR & Friedlander AL (2007) Effects of oligofructose-enriched inulin on intestinal absorption of calcium and magnesium and bone turnover markers in postmenopausal women. *British Journal of Nutrition* **97**, 365-372.

Howard BV, Van HL, Hsia J, *et al.* (2006) Low-fat dietary pattern and risk of cardiovascular disease: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* **295**, 655-666.

Jensen L, Sloth B, Krog-Mikkelsen I, Flint A, Raben A, Tholstrup T, Brunner N & Astrup A (2008) A low-glycemic-index diet reduces plasma plasminogen activator inhibitor-1 activity, but not tissue inhibitor of proteinases-1 or plasminogen activator inhibitor-1 protein, in overweight women. *American Journal of Clinical Nutrition* **87**, 97-105.

Jenkins DJ, Kendall CW, Vuksan V, *et al.* (1999) The effect of wheat bran particle size on laxation and colonic fermentation. *Journal of the American College of Nutrition* **18**, 339-345.

Jenkins DJ, Kendall CW, Vuksan V, Vidgen E, Wong E, Augustin LS & Fulgoni V, III (2000) Effect of cocoa bran on low-density lipoprotein oxidation and fecal bulking. *Archives of Internal Medicine* **160**, 2374-2379.

Ley SJ, Metcalf PA, Scragg RKR & Swinburn BA (2004) Long-term effects of a reduced fat diet intervention on cardiovascular disease risk factors in individuals with glucose intolerance. *Diabetes Res Clin Pract* **63**, 103-112.

Lin J, Zhang SM, Cook NR, Rexrode KM, Liu S, Manson JE, Lee IM & Buring JE (2005) Dietary intakes of fruit, vegetables, and fiber, and risk of colorectal cancer in a prospective cohort of women (United States). *Cancer Causes Control* **16**, 225-233.

Mai V, Flood A, Peters U, Lacey JV, Jr., Schairer C & Schatzkin A (2003) Dietary fibre and risk of colorectal cancer in the Breast Cancer Detection Demonstration Project (BCDDP)

- follow-up cohort. *Int J Epidemiol* **32**, 234-239.
- Maki KC, Rains TM, Kaden VN, Raneri KR & Davidson MH (2007) Effects of a reduced-glycemic-load diet on body weight, body composition, and cardiovascular disease risk markers in overweight and obese adults. *American Journal of Clinical Nutrition* **85**, 724-734.
- McCullough ML, Robertson AS, Chao A, Jacobs EJ, Stampfer MJ, Jacobs DR, Diver WR, Calle EE & Thun MJ (2003) A prospective study of whole grains, fruits, vegetables and colon cancer risk. *Cancer Causes Control* **14**, 959-970.
- McCullough, Marjorie L Robertson, Andrea S Chao, Ann Jacobs, Eric J Stampfer, Meir J Jacobs, David R Diver, William R Calle, Eugenia E Thun, Michael J Netherlands Cancer causes & control : CCC Cancer Causes Control. 2003 Dec;14(10):959-70
- McMillan-Price J, Petocz P, Atkinson F, O'Neill K, Samman S, Steinbeck K, Caterson I & Brand-Miller J (2006) Comparison of 4 diets of varying glycemic load on weight loss and cardiovascular risk reduction in overweight and obese young adults: a randomized controlled trial. *Archives of Internal Medicine* **166**, 1466-1475.
- Michels KB, Fuchs CS, Giovannucci E, Colditz GA, Hunter DJ, Stampfer MJ & Willett WC (2005) Fiber Intake and Incidence of Colorectal Cancer among 76,947 Women and 47,279 Men. *Cancer Epidemiology Biomarkers Prevention* **14**, 842-849.
- Murphy N, Norat T, Ferrari P, *et al.* (2012) Dietary fibre intake and risks of cancers of the colon and rectum in the European prospective investigation into cancer and nutrition (EPIC). *PLoS ONE [Electronic Resource]* **7**.
- Njike VY, Faridi Z, Shuval K, Dutta S, Kay CD, West SG, Kris-Etherton PM & Katz DL (2011) Effects of sugar-sweetened and sugar-free cocoa on endothelial function in overweight adults. *Int J Cardiol* **149**, 83-88.
- Nomura AM, Hankin JH, Henderson BE, Wilkens LR, Murphy SP, Pike MC, Marchand LL, Stram DO, Monroe KR & Kolonel LN (2007) Dietary fiber and colorectal cancer risk: the multiethnic cohort study. *Cancer Causes Control* **18**, 753-764.
- Otani T, Iwasaki M, Ishihara J, Sasazuki S, Inoue M & Tsugane S (2006) Dietary fiber intake and subsequent risk of colorectal cancer: the Japan Public Health Center-based prospective study. *Int J Cancer* **119**, 1475-1480.
- Pelkman CL, Fishell VK, Maddox DH, Pearson TA, Mauger DT & Kris-Etherton PM (2004) Effects of moderate-fat (from monounsaturated fat) and low-fat weight-loss diets on the serum lipid profile in overweight and obese men and women. *American Journal of Clinical Nutrition* **79**, 204-212.
- Pereira MA, Swain J, Goldfine AB, Rifai N & Ludwig DS (2004) Effects of a low-glycemic load diet on resting energy expenditure and heart disease risk factors during weight loss. *JAMA* **292**, 2482-2490.

- Philippou E, McGowan BM, Brynes AE, Dornhorst A, Leeds AR & Frost GS (2008) The effect of a 12-week low glycaemic index diet on heart disease risk factors and 24 h glycaemic response in healthy middle-aged volunteers at risk of heart disease: a pilot study. *European Journal of Clinical Nutrition* **62**, 145-149.
- Philippou E, Bovill-Taylor C, Rajkumar C, Vampa ML, Ntatsaki E, Brynes AE, Hickson M & Frost GS (2009) Preliminary report: the effect of a 6-month dietary glycemic index manipulation in addition to healthy eating advice and weight loss on arterial compliance and 24-hour ambulatory blood pressure in men: a pilot study. *Metabolism* **58**, 1703-1708.
- Pietinen P, Malila N, Virtanen M, Hartman TJ, Tangrea JA, Albanes D & Virtamo J (1999) Diet and risk of colorectal cancer in a cohort of Finnish men. *Cancer Causes Control* **10**, 387-396.
- Poppitt SD, Keogh GF, Prentice AM, Williams DE, Sonnemans HM, Valk EE, Robinson E & Wareham NJ (2002) Long-term effects of ad libitum low-fat, high-carbohydrate diets on body weight and serum lipids in overweight subjects with metabolic syndrome. *American Journal of Clinical Nutrition* **75**, 11-20.
- Raatz SK, Torkelson CJ, Redmon JB, Reck KP, Kwong CA, Swanson JE, Liu C, Thomas W & Bantle JP (2005) Reduced glycemic index and glycemic load diets do not increase the effects of energy restriction on weight loss and insulin sensitivity in obese men and women. *Journal of Nutrition* **135**, 2387-2391.
- Raben A, Vasilaras TH, Moller AC & Astrup A (2002) Sucrose compared with artificial sweeteners: different effects on ad libitum food intake and body weight after 10 wk of supplementation in overweight subjects. *American Journal of Clinical Nutrition* **76**, 721-729.
- Reid M, Hammersley R & Duffy M (2010) Effects of sucrose drinks on macronutrient intake, body weight, and mood state in overweight women over 4 weeks. *Appetite* **55**, 130-136.
- Reid M, Hammersley R, Duffy M & Ballantyne C (2014) Effects on obese women of the sugar sucrose added to the diet over 28 d: a quasi-randomised, single-blind, controlled trial. *Br J Nutr* **111**, 563-570.
- Reid M, Hammersley R, Hill AJ & Skidmore P (2007) Long-term dietary compensation for added sugar: effects of supplementary sucrose drinks over a 4-week period. *British Journal of Nutrition* **97**, 193-203.
- Sanders TA & Reddy S (1992) The influence of rice bran on plasma lipids and lipoproteins in human volunteers. *European Journal of Clinical Nutrition* **46**, 167-172.
- Saris WH, Astrup A, Prentice AM, *et al.* (2000) Randomized controlled trial of changes in dietary carbohydrate/fat ratio and simple vs complex carbohydrates on body weight and blood lipids: the CARMEN study. The Carbohydrate Ratio Management in European National diets. *International Journal of Obesity* **24**, 1310-1318.

- Schatzkin A, Park Y, Leitzmann MF, Hollenbeck AR & Cross AJ (2008) Prospective study of dietary fiber, whole grain foods, and small intestinal cancer. *Gastroenterology* **135**, 1163-1167.
- Shin A, Li H, Shu XO, Yang G, Gao YT & Zheng W (2006) Dietary intake of calcium, fiber and other micronutrients in relation to colorectal cancer risk: Results from the Shanghai Women's Health Study. *Int J Cancer* **119**, 2938-2942.
- Sichieri R, Moura AS, Genelhu V, Hu F & Willett WC (2007) An 18-mo randomized trial of a low-glycemic-index diet and weight change in Brazilian women. *Am J Clin Nutr* **86**, 707-713.
- Tahiri M, Tressol JC, Arnaud J, *et al.* (2003) Effect of short-chain fructooligosaccharides on intestinal calcium absorption and calcium status in postmenopausal women: A stable-isotope study. *American Journal of Clinical Nutrition* **77**, 449-457.
- Tredger JA, Morgan LM, Travis J & Marks V (1991) The effects of guar gum, sugar beet fibre and wheat bran supplementation on serum lipoprotein levels in normocholesterolaemic volunteers. *Journal of Human Nutrition and Dietetics* **4**, 375-384.
- van den Heuvel EG, Muijs T, Brouns F & Hendriks HF (2009) Short-chain fructooligosaccharides improve magnesium absorption in adolescent girls with a low calcium intake. *Nutrition Research* **29**, 229-237.
- van den Heuvel EGHM, Schaafsma G, Muys T & Van Dokkum W (1998) Nondigestible oligosaccharides do not interfere with calcium and nonheme-iron absorption in young, healthy men. *American Journal of Clinical Nutrition* **67**, 445-451.
- van den Heuvel EGHM, Schoterman MHC & Muijs T (2000) Transgalactooligosaccharides Stimulate Calcium Absorption in Postmenopausal Women. *J Nutr* **130**, 2938-2942.
- Wakai K, Date C, Fukui M, *et al.* (2007) Dietary Fiber and Risk of Colorectal Cancer in the Japan Collaborative Cohort Study. *Cancer Epidemiology Biomarkers Prevention* **16**, 668-675.
- Wolever TM & Mehling C (2003) Long-term effect of varying the source or amount of dietary carbohydrate on postprandial plasma glucose, insulin, triacylglycerol, and free fatty acid concentrations in subjects with impaired glucose tolerance. *American Journal of Clinical Nutrition* **77**, 612-621.
- Vuksan V, Jenkins DJ, Vidgen E, Ransom TP, Ng MK, Culhane CT & O'Connor D (1999) A novel source of wheat fiber and protein: effects on fecal bulk and serum lipids. *American Journal of Clinical Nutrition* **69**, 226-230.