

# 19

## Influenza

### The disease

Influenza is an acute viral infection of the respiratory tract. There are three types of influenza virus: A, B and C. Influenza A and influenza B are responsible for most clinical illness. Influenza is highly infectious with a usual incubation period of one to three days.

The disease is characterised by the sudden onset of fever, chills, headache, myalgia and extreme fatigue. Other common symptoms include a dry cough, sore throat and stuffy nose. For otherwise healthy individuals, influenza is an unpleasant but usually self-limiting disease with recovery usually within two to seven days. The illness may be complicated by (and may present as) bronchitis, secondary bacterial pneumonia or, in children, otitis media. Influenza can be complicated more unusually by meningitis, encephalitis or meningoencephalitis. The risk of serious illness from influenza is higher amongst children under six months of age (Poehling *et al.*, 2006; Ampfo *et al.*, 2006; Coffin *et al.*, 2007; Zhou *et al.*, 2012), older people (Thompson *et al.*, 2003 and 2004; Zhou *et al.*, 2012) and those with underlying health conditions such as respiratory or cardiac disease, chronic neurological conditions, or immunosuppression and pregnant women (Neuzil *et al.*, 1998; O'Brien *et al.*, 2004; Nicoll *et al.*, 2008 and Pebody *et al.*, 2010). Influenza during pregnancy may also be associated with perinatal mortality, prematurity, smaller neonatal size and lower birth weight (Pierce *et al.*, 2011; Mendez-Figueroa *et al.*, 2011). Although primary influenza pneumonia is a rare complication that may occur at any age and carries a high case fatality rate (Barker and Mullooly, 1982), it was seen more frequently during the 2009 pandemic and the following influenza season. Serological studies in healthcare professionals have shown that approximately 30 to 50% of influenza infections can be asymptomatic (Wilde *et al.*, 1999) but the proportion of influenza infections that are asymptomatic may vary depending on the characteristics of the influenza strain.

Transmission is by droplets, aerosol, or through direct contact with respiratory secretions of someone with the infection (Lau *et al.*, 2010). Influenza spreads

rapidly, especially in closed communities. Most cases in the UK tend to occur during an eight- to ten-week period during the winter. The timing, extent and severity of this ‘seasonal’ influenza can all vary. Influenza A viruses cause outbreaks most years and it is these viruses that are the usual cause of epidemics. Large epidemics occur intermittently. Influenza B tends to cause less severe disease and smaller outbreaks overall. The burden of influenza B disease is mostly in children when the severity of illness can be similar to that associated with influenza A.

Changes in the principal surface antigens of influenza A – haemagglutinin and neuraminidase – make these viruses antigenically labile. Minor changes described as antigenic drift occur progressively from season to season. Antigenic shift occurs periodically, resulting in major changes and the emergence of a new subtype with a different haemagglutinin protein. Because immunity from the previous virus may not protect completely against the new subtype, the population may have little or no immunity, and this may therefore lead to widespread epidemics or even a pandemic.

Three influenza pandemics occurred in the last century (in 1918, 1957 and 1968). The first influenza pandemic of this century was declared by the World Health Organization (WHO) in June 2009. This was caused by an influenza A(H1N1)v virus. The influenza A(H1N1)v pandemic caused higher rates of illness in children and young adults and lower rates of illness in adults aged 60 years and older when compared with ‘seasonal’ influenza (Writing Committee of the WHO Consultation on Clinical Aspects of Pandemic (H1N1) 2009 Influenza, 2010). For most individuals the disease was mild. Symptoms were similar to those of ‘seasonal’ influenza, although gastrointestinal symptoms (vomiting and diarrhoea) were more commonly reported than is usual for ‘seasonal’ influenza. During the pandemic, there were fewer than 500 laboratory confirmed deaths from influenza A(H1N1)v in the UK with an overall estimated case fatality ratio of 0.25 per 1000 clinical cases (95% confidence limits 0.13 to 0.4 per 1000 clinical cases) (Presanis, *et al.*, 2011). The highest mortality rates were in those with chronic neurological disease, respiratory disease and immunosuppression (Pebody *et al.*, 2010). Individuals with morbid obesity (BMI>40) were also found to be at higher risk of severe outcome (both hospitalisation and death) following pandemic influenza infection compared to individuals with obesity and to normal weight individuals (Morgan *et al.*, 2010; Fezeu *et al.*, 2011; Van Kerkhove, 2011). Pregnant women were also at increased risk of complications (Jamison *et al.*, 2009). Most of the serious complications arising from influenza A(H1N1) v infection occurred in people with underlying health conditions, but a significant proportion arose

in people who had been previously healthy (Writing Committee of the WHO Consultation on Clinical Aspects of Pandemic (H1N1) 2009 Influenza, 2010).




Figure 19.1 Rate of influenza/influenza-like illness episodes in England (weekly returns to Royal College of General Practitioners), 2009–10 to 2014–15 showing the variation in the timing and shape of influenza activity usually between weeks 37 and 15. However, much of the influenza A(H1N1)v pandemic activity was outside that usual time window, demonstrated by the non-seasonal peak at week 29 in 2009/10. Data for 2009/10 may underestimate the extent of influenza-like illness due to the introduction of the National Pandemic Flu Service in England during 2009. There may be differences in the epidemiology of influenza between the different countries in the UK. Data provided by PHE (formerly HPA) and RCGP.

The influenza A(H1N1)v strain continued to cause widespread illness during the 2010/11 influenza season. Despite the recent emergence of the influenza A(H1N1)v strain, conditions still exist for the emergence of future influenza strains with potential to lead to another pandemic (for example, from influenza A H5N1 or H7N9 strains).

Influenza B viruses are also subject to antigenic drift but with less frequent changes.

## History and epidemiology of the disease

Influenza activity is monitored in the UK through a variety of schemes based in primary and secondary care. One important indicator is based on reports

of new consultations for influenza-like illness from sentinel GP practices, combined with virological surveillance. Weekly reports are collated by Public Health England (PHE, formerly Health Protection Agency (HPA)). Additional information for England is provided by the Royal College of General Practitioners (RCGP), for Scotland by Health Protection Scotland, for Wales by Public Health Wales and for Northern Ireland by the Public Health Agency.

Official estimates of the number of deaths attributable to influenza are produced by PHE. These are inferred from the number of all-cause death registrations in the winter period that are above an expected seasonal level. However, as the cause of deaths is not examined directly, deaths above the expected level may include causes other than influenza such as cold weather related conditions. Estimates of excess winter deaths potentially attributable to influenza in years in the last decade in England and Wales range from not determined (in 2005-6 and 2006-7) to 10,351 (in 2008-9). The highest estimate in the past two decades was 21,497 for the 1999-2000 influenza season (Donaldson *et al.*, 2010).

PHE also collects data on admissions to intensive care units and on deaths with a laboratory-confirmed influenza infection. Whilst it is not possible to ascertain all fatal cases where influenza was involved, investigation of such cases allows assessment of the characteristics of people most severely affected by influenza, including age and the responsible influenza type. An analysis by PHE of data from fatal cases collected in England during the 2010/11 influenza season, when influenza A(H1N1)v was the predominant circulating strain, gives an indication of the increased risk of death from influenza complications for those in clinical risk groups (see Table 19.1).

Table 19.1 Influenza-related population mortality rates and relative risk of death among those aged six months to under 65 years by clinical risk group in England, September 2010 – May 2011.

	Number of fatal flu cases (%)	Mortality rate per 100,000 population	Age-adjusted relative risk*
<b>In a risk group</b>	213 (59.8)	4.0	11.3 (9.1-14.0)
<b>Not in any risk group</b>	143 (40.2)	0.4	Baseline
<b>Chronic renal disease</b>	19 (5.3)	4.8	18.5

	Number of fatal flu cases (%)	Mortality rate per 100,000 population	Age-adjusted relative risk*
<b>Chronic heart disease</b>	32 (9.0)	3.7	10.7 (7.3-15.7)
<b>Chronic respiratory disease</b>	59 (16.6)	2.4	7.4 (5.5-10.0)
<b>Chronic liver disease</b>	32 (9.0)	15.8	48.2 (32.8-70.6)
<b>Diabetes</b>	26 (7.3)	2.2	5.8 (3.8-8.9)
<b>Immunosuppression</b>	71 (19.9)	20.0	47.3 (35.5-63.1)
<b>Chronic neurological disease (excluding stroke/transient ischaemic attack)</b>	42 (11.8)	14.7	40.4 (28.7-56.8)
<b>Total (including 22 cases with no information on clinical risk factors)</b>	378	0.8	

\* Mantel-Haenszel age-adjusted rate ratio (RR), with corresponding exact 95% CI were calculated for each risk group using the two available age groups (from six months up to 15 years and from 16 to 64 years).

Table reproduced from *Surveillance of influenza and other respiratory viruses in the UK 2010-2011 report* by kind permission of PHE.

Influenza immunisation has been recommended in the UK since the late 1960s, with the aim of directly protecting those in clinical risk groups who are at a higher risk of influenza associated morbidity and mortality. In 2000, the policy was extended to include all people aged 65 years or over (see later for age definition). The list of conditions that constitute a clinical risk group where influenza vaccine is indicated are reviewed regularly by the Joint Committee on Vaccination and Immunisation (JCVI). In 2010, pregnancy was added as a clinical risk category, and in October 2014 the JCVI advised that morbid

obesity (defined as BMI 40+) should be considered a risk factor for seasonal influenza vaccination.

Uptake of influenza vaccination in those aged 65 years or over and in those aged under 65 years in a clinical risk group (excluding data on pregnant women) in the UK is shown in Table 19.2.

Table 19.2 Influenza vaccine uptake in the UK for people aged 65 years or over and, in brackets, aged under 65 years in a clinical risk group (excluding pregnant women). End of influenza vaccination campaign estimates.

Year	England (%)	Scotland (%)	Wales (%)	Northern Ireland (%)
2000–01	65.4	65	39	68
2001–02	67.5	65	59	72
2002–03	68.6	69	54	72.1 (55.8)
2003–04	71.0	72.5	63	73.4 (63.8)
2004–05	71.5 (39.9)	71.7 (39.3)	63	72.7 (65.2)
2005–06	75.3 (48.0)	77.8 (46.3)	68	76.8 (80.9)
2006–07	73.9 (42.1)	75.2 (37.8)	*	75.1 (71.2)
2007–08	73.5 (45.3)	74.3 (44.4)	64	75.7 (68.3)
2008–09	74.1 (47.1)	76.3 (47.8)	60 (41)	76.8 (74.0)
2009–10	72.4 (51.6)	75.0 (53.4)	64 (49)	77.0 (80.0)
2010–11	72.8 (50.4)	75.3 (56.1)	65.8 (48.6)	74.9 (78.7)
2011–12	74.0 (51.6)	76.6 (59.7)	67.7 (50.0)	77.0 (81.7)
2012–13	73.4 (51.3)	77.4 (59.2)	67.7 (49.7)	75.0 (80.2)
2013–14	73.2 (52.3)	76.9% (57.5) **	68.3 (51.1)	75.4 (76.4)
2014–15	72.8 (50.3)**	76.3 (54.0)**	68.1 (49.5)**	73.4 (71.8)

\* Data not available. \*\*Provisional data.

The uptake of influenza vaccine by pregnant women is difficult to estimate as it is more challenging to determine a denominator accurately. The available data are shown in Table 19.3 but may underestimate uptake.

Table 19.3 Influenza vaccine uptake in the UK since the start of the influenza immunisation programme for pregnant women. End of influenza vaccination campaign estimates.

Year	England (%)	Scotland (%)	Wales (%)	Northern Ireland (%)
2010-11	38.0	65.6*	39.6	N/A
2011-12	27.4	41.1	31.7	58.4
2012-13	40.3	54.0	61.6	64.6
2013-14	39.8	49.2	43.8	58.0
2014-15	44.1	50.9	45.3**†	56.1

\* Denominator incomplete \*\*Provisional data †GP practice data, post natal survey uptake 72.4%

Information on vaccine uptake for the childhood programme and more detailed analyses of influenza vaccine uptake by individual clinical risk groups and by different age groups are made available by the UK public health bodies on their webpages.

### Extension of the influenza programme to Children

In 2012, JCVI recommended that the programme should be extended to all children aged two to less than seventeen years old (JCVI, 2012). JCVI advised that the vaccine of choice for the extension to the programme should be live attenuated intranasal influenza vaccine (LAIV Fluenz Tetra®), given the evidence of superior effectiveness in young children, particularly after a single dose, and the potential protection against drifted strains. The route of administration also makes LAIV an easier vaccine to administer and more acceptable to parents and children when compared to an injectable vaccine.

The phased introduction of the extension of the influenza programme to children began in 2013. Those cohorts eligible for the programme in each UK country will be updated each season in the annual flu letter for England and the respective Chief Medical Officer letters for the Devolved Administrations.

England:

<https://www.gov.uk/government/collections/annual-flu-programme>

Northern Ireland:

[http://www.dhsspsni.gov.uk/index/phealth/professional/cmo\\_communications.htm](http://www.dhsspsni.gov.uk/index/phealth/professional/cmo_communications.htm)

Scotland:

<http://www.sehd.scot.nhs.uk/index.asp?name=&org=%25&keyword=&category=9&number=10&sort=tDate&order=DESC&Submit=Go>

Wales:

<http://gov.wales/topics/health/cmo/publications/cmo/?lang=en>

## The influenza vaccination

All but one of the influenza vaccines available in the UK are inactivated and do not contain live viruses. One vaccine (Fluenz Tetra<sup>®</sup>) contains live viruses that have been attenuated (weakened) and adapted to cold so that they cannot replicate efficiently at body temperature. None of the influenza vaccines can therefore cause clinical influenza in those that can be vaccinated, although mild coryzal symptoms can occur with the live vaccine. The live vaccine (Fluenz Tetra<sup>®</sup>) is administered by nasal spray, and the inactivated vaccines are all administered by intramuscular injection with the exception of one preparation (Intanza<sup>®</sup>) which is administered by the intradermal route. Most of the vaccines are prepared from viruses grown in embryonated hens eggs.

Because of the changing nature of influenza viruses, WHO monitors the epidemiology of influenza viruses throughout the world. Each year it makes recommendations about the strains to be included in vaccines for the forthcoming winter for the northern and southern hemispheres (<http://www.who.int/influenza/en/>).

Influenza vaccines are prepared using virus strains in line with the WHO recommendations. Most current inactivated influenza vaccines are trivalent, containing two subtypes of influenza A and one B virus; however, quadrivalent vaccines with an additional B virus have been developed. The first authorised quadrivalent influenza vaccine became available for use in the UK in 2013.

Manufacture of influenza vaccines is complex and conducted to a tight schedule, constrained by the period between the announcement of the WHO recommendations and the opportunity to vaccinate before the influenza season. Manufacturers may not be able to respond to unexpected demands for vaccine at short notice.

With the exception of 2014/15, in most recent years, the vaccines have closely matched the influenza A viruses circulating during the influenza season. In

the northern hemisphere in 2014/15, however, antigenic drift in the A(H3N2) viruses was observed such that the H3N2 component of the vaccine did not provide optimal protection. Mismatches between the components in the vaccine and circulating viruses do occur from time to time and explains much of the variation in estimates of vaccine effectiveness (Osterholm *et al.*, 2012). When antigenic drift does occur, vaccination is still recommended as some degree of protection may be conferred against the drifted strain and the vaccine should still offer protection against the other strains in the vaccine. The live attenuated influenza vaccine is also thought to provide broader protection than inactivated vaccines, and therefore has potential to offer better protection against strains that have undergone antigenic drift compared to the original virus strains in the vaccine (Ambose *et al.*, 2011; Hoft *et al* 2011; Subbramanian *et al* 2010).

Because the trivalent vaccine contains an influenza B strain from a single lineage, mismatches between the vaccine and the circulating B strain occur more frequently. The use of quadrivalent influenza vaccines containing a B strain from each lineage is expected to improve the matching of the vaccine in the future.

JCVI has advised that, all other things being equal, quadrivalent inactivated vaccine is preferable to trivalent inactivated influenza vaccine. For adults in clinical risk groups a range of inactivated vaccines are available from suppliers for GPs to purchase. Because influenza B is relatively more common in children, the vaccines centrally purchased for the childhood programme in recent years have been quadrivalent preparations (Fluenz Tetra<sup>®</sup> and Fluarix Tetra<sup>®</sup>). The childhood programme should therefore contribute to better control of influenza B overall, by reducing transmission across the population.

If a new influenza A subtype were to emerge with epidemic or pandemic potential (as occurred in 2009 with influenza A(H1N1)v), it is unlikely that the influenza vaccine will be well matched to the emerging strain. In these circumstances, as occurred during the second wave of the 2009 pandemic, a monovalent vaccine against that strain may be developed and implemented.

All authorised influenza vaccines need to meet immunogenicity, safety and quality criteria set by the European Committee for Medicinal Products for Human Use (CHMP), with the assessment of expected efficacy based on meeting or exceeding indicated requirements in serological assessments of immunogenicity (EMA, 1997). A recent meta-analysis, which included studies when the influenza virus strains in the trivalent vaccine were drifted or mismatched with those in circulation, suggested an overall effectiveness against

confirmed disease of 59% (95% confidence interval 51-67) in adults aged 18 to 65 years (Osterholm *et al.*, 2012). In the elderly, protection produced by the vaccine may be lower (Fleming *et al.*, 2010), although immunisation has been shown to reduce the incidence of severe disease including bronchopneumonia, hospital admissions and mortality (Wright *et al.*, 1977; Mangtani *et al.*, 2004). Live attenuated influenza vaccine has been shown to provide a higher level of protection for children than trivalent inactivated influenza vaccine (Belshe *et al.*, 2007); a recent meta-analysis suggested an efficacy against confirmed disease of 83% (95% confidence interval 69-91) (Osterholm *et al.*, 2012; Ashkenazi *et al.*, 2006; Fleming *et al.*, 2006).

After immunisation, protective immune responses may be achieved within 14 days. Although influenza activity is not usually significant in the UK before the middle of November, the influenza season can start early (as it did in 2003–04), and therefore the ideal time for immunisation is between September and early November. Protection afforded by the vaccine is thought to last for at least one influenza season. However, as the level of protection provided in subsequent seasons is likely to reduce and there may be changes to the circulating strains from one season to the next, annual revaccination is important.

A list of the influenza vaccines available in the UK is published ahead of the influenza season in the annual flu letter for England (available at: <https://www.gov.uk/government/collections/annual-flu-programme>).

### Storage (also refer to Chapter 3)

Vaccines should be stored in the original packaging at +2°C to +8°C and protected from light. All vaccines are sensitive to some extent to heat and cold. Heat speeds up the decline in potency of most vaccines, thus reducing their shelf life. Efficacy, safety and quality may be adversely affected if vaccines are not stored at the temperatures specified in the license. Freezing may cause increased reactogenicity and loss of potency for some vaccines and can also cause hairline cracks in the container, leading to contamination of the contents.

Fluenz Tetra® may be left out of the refrigerator for a maximum period of 12 hours at a temperature not above 25°C as indicated in the SPC. If the vaccine has not been used after this 12 hour period, it should be disposed of.

### Presentation

Inactivated influenza vaccines for intramuscular administration are supplied as suspensions in pre-filled syringes. They should be shaken well before they are administered.

Intanza<sup>®</sup>, the intradermal vaccine, is supplied in a micro-needle injection system.

Fluenz Tetra<sup>®</sup>, the intranasal vaccine, is supplied as a nasal spray suspension in a special applicator.

## Dosage and schedule

The dosages and schedules for influenza vaccines should be given according to the recommendations for use of the vaccines (see later).

Some of the summaries of product characteristics (SPCs) for intramuscular inactivated influenza vaccines indicate that young children can be given either a 0.25ml or a 0.5ml dose. JCVI has advised that where these alternative doses are indicated in the SPC, the 0.5ml dose of intramuscular inactivated influenza vaccine should be given to infants aged six months or older and young children because there is evidence that this dose is effective in young children (Heinonen *et al.*, 2011).

Children aged six months to under nine years who are in clinical risk groups and have not received influenza vaccine previously should be offered a second dose of vaccine. JCVI has advised that children aged two years to under nine years of age who are not in a clinical risk group, only require a single dose of live attenuated influenza vaccine irrespective of whether they have received influenza vaccine previously. This advice differs from that in the SPC for Fluenz Tetra. Children who have received one or more doses of any influenza vaccine before (including pandemic monovalent influenza A(H1N1)v vaccine) should be considered as previously vaccinated (see later section on children).

## Administration

The inactivated influenza vaccines given by intramuscular injection should be given preferably into the upper arm (or anterolateral thigh in infants). However, individuals with a bleeding disorder should be given vaccine by deep subcutaneous injection to reduce the risk of bleeding.

The inactivated influenza vaccine administered by the intradermal route (Intanza<sup>®</sup>) is supplied in a micro-needle injection system that is held at right-angles to the skin. The device allows intradermal vaccination to be performed without the need for additional training.

The live attenuated influenza vaccine is administered by the intranasal route (Fluenz Tetra<sup>®</sup>) and is supplied in an applicator that allows a divided dose to

be administered in each nostril (total dose of 0.2ml, 0.1ml in each nostril). The device allows intranasal administration to be performed without the need for additional training. Administration of either dose does not need to be repeated if the patient sneezes or blows their nose following administration. There are no data on the effectiveness of Fluenz Tetra® when given to children with a heavily blocked or runny nose (rhinitis) attributable to infection or allergy. As heavy nasal congestion might impede delivery of the vaccine to the nasopharyngeal mucosa, deferral of administration until resolution of the nasal congestion or an appropriate alternative intramuscularly administered influenza vaccine should be considered.

Inactivated influenza vaccines can be given at the same time as other vaccines. The live attenuated vaccine can also be given at the same time as other live or inactivated vaccines. Although it was previously recommended that, where vaccines cannot be administered simultaneously; a four-week interval should be observed between live viral vaccines, JCVI have advised that no specific intervals need to be observed between the live attenuated intranasal influenza vaccine and other live vaccines (see [Chapter 6](#)). Intramuscular and intradermal vaccines should be given at separate sites, preferably in a different limb. If given in the same limb, they should be given at least 2.5cm apart (American Academy of Pediatrics, 2003).

As a wide variety of influenza vaccines are on the UK market each year, it is especially important that the exact brand of vaccine, batch number and site at which each vaccine is given is accurately recorded in the patient records. Where the vaccine is given for occupational reasons, it is recommended that the employer keep a vaccination record. It is important that vaccinations given either at a general practice or elsewhere (for example, at community pharmacies, or antenatal clinics) are recorded on appropriate health records for the individual (using the appropriate clinical code) in a timely manner, including the relevant Child Health Information System. If given elsewhere, a record of vaccination should be returned to the patient's general practice to allow clinical follow up and to avoid duplicate vaccination.

## Recommendations for the use of the vaccines

The objectives of the influenza immunisation programme are to protect those who are most at risk of serious illness or death should they develop influenza and to reduce transmission of the infection, thereby contributing to the protection of vulnerable patients who may have a suboptimal response to their own immunisations.

To facilitate this, general practitioners are required to proactively identify all those for whom influenza immunisations are indicated and to compile a register of those patients for whom influenza immunisation is recommended. Sufficient vaccine can then be ordered in advance and patients can be invited to planned immunisation sessions or appointments. Given that some influenza vaccines are restricted for use in particular age groups, the SPCs for individual products should always be referred to when ordering vaccines to ensure that they can be given appropriately to particular patient age groups.

Research has identified processes at GP surgeries that are associated with higher uptake of influenza vaccine (Dexter *et al.*, 2012). This included, having a named individual at the surgery responsible for the influenza immunisation programme; update and maintenance of an accurate register of patients eligible for influenza immunisation and direct contact with eligible patients inviting them for immunisation.

Patients should be advised that many other organisms cause respiratory infections similar to influenza during the influenza season, e.g. the common cold and respiratory syncytial virus. Influenza vaccine will not protect against these diseases.

Influenza vaccine should be offered, ideally before influenza viruses start to circulate, to:

- all those aged 65 years or older (for definition please see the annual flu letter for the coming/current season)
- all those aged six months or older in the clinical risk groups shown in Table 19.4
- children not in clinical risk groups that are eligible for vaccination as part of the ongoing phased roll out of the extension of the programme to all children aged two to less than seventeen years old.\*

\* Note: please see the respective annual flu letters for England and the Devolved Administrations for specific details on the cohorts of children that are eligible for influenza vaccination.

Table 19.4 Clinical risk groups who should receive the influenza immunisation. Influenza vaccine should be offered to people in the clinical risk categories set out below.

Clinical risk category	Examples (this list is not exhaustive and decisions should be based on clinical judgement)
<b>Chronic respiratory disease</b>	<p>Asthma that requires continuous or repeated use of inhaled or systemic steroids or with previous exacerbations requiring hospital admission.</p> <p>Chronic obstructive pulmonary disease (COPD) including chronic bronchitis and emphysema; bronchiectasis, cystic fibrosis, interstitial lung fibrosis, pneumoconiosis and bronchopulmonary dysplasia (BPD).</p> <p>Children who have previously been admitted to hospital for lower respiratory tract disease.</p> <p><i>see precautions section on live attenuated influenza vaccine</i></p>
<b>Chronic heart disease</b>	<p>Congenital heart disease, hypertension with cardiac complications, chronic heart failure, individuals requiring regular medication and/or follow-up for ischaemic heart disease.</p>
<b>Chronic kidney disease</b>	<p>Chronic kidney disease at stage 3, 4 or 5, chronic kidney failure, nephrotic syndrome, kidney transplantation.</p>
<b>Chronic liver disease</b>	<p>Cirrhosis, biliary atresia, chronic hepatitis</p>
<b>Chronic neurological disease (included in the DES directions for Wales)</b>	<p>Stroke, transient ischaemic attack (TIA). Conditions in which respiratory function may be compromised due to neurological disease (e.g. polio syndrome sufferers). Clinicians should offer immunisation, based on individual assessment, to clinically vulnerable individuals including those with cerebral palsy, learning disabilities, multiple sclerosis and related or similar conditions; or hereditary and degenerative disease of the nervous system or muscles; or severe neurological disability.</p>
<b>Diabetes</b>	<p>Type 1 diabetes, type 2 diabetes requiring insulin or oral hypoglycaemic drugs, diet controlled diabetes.</p>

Clinical risk category	Examples (this list is not exhaustive and decisions should be based on clinical judgement)
<p><b>Immunosuppression</b> (see <i>contraindications and precautions section on live attenuated influenza vaccine</i>)</p>	<p>Immunosuppression due to disease or treatment, including patients undergoing chemotherapy leading to immunosuppression, bone marrow transplant, HIV infection at all stages, multiple myeloma or genetic disorders affecting the immune system (e.g. IRAK-4, NEMO, complement disorder)</p> <p>Individuals treated with or likely to be treated with systemic steroids for more than a month at a dose equivalent to prednisolone at 20mg or more per day (any age), or for children under 20kg, a dose of 1mg or more per kg per day.</p> <p>It is difficult to define at what level of immunosuppression a patient could be considered to be at a greater risk of the serious consequences of influenza and should be offered influenza vaccination. This decision is best made on an individual basis and left to the patient's clinician.</p> <p>Some immunocompromised patients may have a suboptimal immunological response to the vaccine.</p>
<p><b>Asplenia or dysfunction of the spleen</b></p>	<p>This also includes conditions such as homozygous sickle cell disease and coeliac syndrome that may lead to splenic dysfunction.</p>
<p><b>Pregnant women</b></p>	<p>Pregnant women at any stage of pregnancy (first, second or third trimesters).</p> <p><i>see precautions section on live attenuated influenza vaccine</i></p>
<p><b>Morbid obesity (class III obesity)*</b></p>	<p>Adults with a Body Mass Index <math>\geq 40</math> kg/m<sup>2</sup></p>

\* Many of this patient group will already be eligible due to complications of obesity that place them in another risk category

### Other groups

The list above is not exhaustive, and the medical practitioner should apply

clinical judgment to take into account the risk of influenza exacerbating any underlying disease that a patient may have, as well as the risk of serious illness from influenza itself. Influenza vaccine should be offered in such cases even if the individual is not in the clinical risk groups specified above. Vaccination should also be offered to household contacts of immunocompromised individuals, i.e. individuals who expect to share living accommodation on most days over the winter and therefore for whom continuing close contact is unavoidable. This may include carers (see below).

In addition to the above, immunisation should be provided to healthcare and social care workers in direct contact with patients/clients to protect them and to reduce the transmission of influenza within health and social care premises, to contribute to the protection of individuals who may have a suboptimal response to their own immunisations, and to avoid disruption to services that provide their care. This would include:

- health and social care staff directly involved in the care of their patients or clients
- those living in long-stay residential care homes or other long-stay care facilities where rapid spread is likely to follow introduction of infection and cause high morbidity and mortality (this does not include prisons, young offender institutions, university halls of residence etc.)
- those who are in receipt of a carer's allowance, or those who are the main carer of an elderly or disabled person whose welfare may be at risk if the carer falls ill. Vaccination should be given on an individual basis at the GP's discretion in the context of other clinical risk groups in their practice
- others involved directly in delivering health and social care such that they and vulnerable patients/clients are at increased risk of exposure to influenza (further information is provided in guidance from UK health departments).

## Children

Studies suggest that two doses of inactivated influenza vaccine may be required to achieve adequate antibody levels in younger children who have not received influenza vaccine before (Allison *et al.*, 2006; Neuzil *et al.*, 2006; Ritzwoller *et al.*, 2005; Shuler *et al.*, 2007; Wright *et al.*, 1977). Live attenuated influenza vaccine has been shown to provide greater protection for children than inactivated influenza vaccine (Belshe *et al.*, 2007; Ashkenazi *et al.*, 2006; Fleming *et al.*, 2006) and studies have also shown meaningful efficacy after a single dose of live attenuated influenza vaccine in previously unvaccinated children (Bracco Neto *et al.*, 2009; Block *et al.*, 2009). Given this, JCVI has advised, as set out below, the use of different dosage schedules of influenza

vaccine for children depending on their age, the clinical indications, the type of vaccine offered and whether they have received influenza vaccine previously. This advice differs from some of the SPCs.

### **Children aged two to less than seventeen years old NOT IN clinical risk groups**

Starting from September 2013, an extension of the programme to all children aged two to less than seventeen years old is being phased in from the youngest age groups. Please see the respective annual flu letters for England and the Devolved Administrations for the cohorts of children that are eligible for influenza vaccination for the coming/current season.

A single dose of Fluenz Tetra® should be offered, irrespective of whether influenza vaccine has been received previously.

### **Children aged six months to less than two years of age IN clinical risk groups**

These children should be offered inactivated trivalent influenza vaccine. Those who have not received influenza vaccine previously should be offered a second dose of vaccine, at least four weeks later. The inactivated influenza vaccines are interchangeable; the second dose, if required, should be given at least four weeks after the first dose in accordance with the manufacturer's SPC for that vaccine.

### **Children aged two to less than 18 years of age IN clinical risk groups**

Children aged two years to less than 18 years in clinical risk groups should be offered Fluenz Tetra® unless it is medically contraindicated (see contraindications and precautions sections). Those children who have never received influenza vaccine before and are aged between two and less than nine years should be offered a second dose of Fluenz Tetra® at least four weeks later. If Fluenz Tetra is unavailable for this second dose (due to batch expiry) an inactivated influenza vaccine can be given.

For those children for whom Fluenz Tetra® is medically contraindicated, a suitable inactivated influenza vaccine should be offered. The quadrivalent inactivated influenza vaccine (Fluarix™ Tetra) is authorised for children from the age of three years and is preferred because of the additional protection offered. The quadrivalent vaccine has both lineages of influenza B and may therefore provide better protection against the circulating B strain(s) than trivalent inactivated influenza vaccines. Otherwise, children aged two years should be given an inactivated trivalent vaccine. Children aged two to less than nine years old offered inactivated influenza vaccine who have not received

influenza vaccine previously should be offered a second dose of the vaccine at least four weeks later.

The inactivated influenza vaccines are interchangeable; the second dose, if required, should be given at least four weeks after the first dose in accordance with the manufacturer's SPC for that vaccine.

Table 19.5 summarises the advice on influenza vaccination for children

## Preterm infants

It is important that preterm infants who have risk factors have their immunisations at the appropriate chronological age. Influenza immunisation should be considered after the child has reached six months of age.

Table 19.5 Influenza vaccination for children under 18 years old

Eligible cohort	Vaccine available: Children in clinical risk groups*	Vaccine available: Children not in clinical risk groups <sup>1</sup>
Six months to less than two years old	Offer suitable inactivated flu vaccine.	Not applicable.
Children aged two years to less than 18 years old <sup>1</sup>	Offer LAIV (Fluenz Tetra <sup>®</sup> ) (unless medically contraindicated <sup>2</sup> )	Offer LAIV (Fluenz Tetra <sup>®</sup> )

\* Children in clinical risk groups aged two to less than nine years who have not received flu vaccine before should be offered two doses of Fluenz Tetra<sup>®</sup> or inactivated vaccine (given at least four weeks apart).

1 Please see the respective annual flu letters for England and the Devolved Administrations for the cohorts of children not in clinical risk groups that are eligible for influenza vaccination for the coming/current season.

2 If Fluenz Tetra<sup>®</sup> is medically contraindicated or otherwise unsuitable, then offer suitable inactivated flu vaccine. If quadrivalent inactivated vaccine available, consider for children age 3 years and older only. Children aged two to less than 9 years who have not received flu vaccine before require two doses of inactivated vaccine (given at least four weeks apart).

## Pregnancy

Pregnant women should be offered inactivated influenza vaccine as the risk of serious illness from influenza is higher in pregnant women (Pebody *et al.*, 2010). In addition, a number of studies show that influenza vaccination during pregnancy provides passive immunity against influenza to infants in the first few months of life following birth (Benowitz *et al.*, 2010; Eick *et al.*,

2010; Zaman et al., 2008; Poehling et al., 2011). A study showed that influenza vaccination reduced the likelihood of prematurity and smaller infant size at birth associated with influenza infection (Omer et al., 2011).

A review of studies on the safety of influenza vaccine in pregnancy concluded that inactivated influenza vaccine can be safely and effectively administered during any trimester of pregnancy and that no study to date has demonstrated an increased risk of either maternal complications or adverse foetal outcomes associated with inactivated influenza vaccine (Tamma et al., 2009). Data are more limited for the live attenuated influenza vaccine (Fluenz Tetra®). Whilst there is no evidence of risk with live attenuated influenza vaccine (Toback et al., 2012), inactivated influenza vaccines are preferred for those who are pregnant. There is no need, however, to specifically test eligible girls for pregnancy or to advise avoidance of pregnancy in those who have been recently vaccinated.

## Contraindications

The SPCs for individual products should always be referred to when deciding which vaccine to give. There are very few individuals who cannot receive any influenza vaccine. When there is doubt, appropriate advice should be sought promptly from the screening and immunisation team in the NHS England area team, a consultant in communicable disease control or a consultant paediatrician, so that the period the individual is left unvaccinated is minimised.

None of the influenza vaccines should be given to those who have had:

- a confirmed anaphylactic reaction to a previous dose of the vaccine, or
- a confirmed anaphylactic reaction to any component of the vaccine (other than ovalbumin – see precautions).

Confirmed anaphylaxis is rare (see Chapter 8 for further information). Other allergic conditions such as rashes may occur more commonly and are not contraindications to further immunisation. A careful history of the event will often distinguish between true anaphylaxis and other events that are either not due to the vaccine or are not life threatening. In the latter circumstance, it may be possible to continue the immunisation course. Specialist advice must be sought on the vaccines and the circumstances in which they could be given (see Chapter 6 for further information). The risk to the individual of not being immunised must be taken into account.

The live attenuated influenza vaccine (Fluenz Tetra®) should not be given to children or adolescents who are clinically severely immunodeficient due

to conditions or immunosuppressive therapy such as: acute and chronic leukaemias; lymphoma; HIV infection not on highly active antiretroviral therapy (HAART); cellular immune deficiencies; and high dose corticosteroids. It is not contraindicated for use in children or adolescents with stable HIV infection receiving antiretroviral therapy; or who are receiving topical corticosteroids, standard dose inhaled corticosteroids or low-dose systemic corticosteroids, or those receiving corticosteroids as replacement therapy, e.g. for adrenal insufficiency. It is contraindicated in children and adolescents receiving salicylate therapy (other than for topical treatment of localised conditions) because of the association of Reye's syndrome with salicylates and wild-type influenza infection as described in the SPC for Fluenz Tetra®.

## Precautions

Minor illnesses without fever or systemic upset are not valid reasons to postpone immunisation. If an individual is acutely unwell, immunisation may be postponed until they have fully recovered. This is to avoid confusing the differential diagnosis of any acute illness by wrongly attributing any signs or symptoms to the adverse effects of the vaccine.

## Immunosuppression and HIV infection

Individuals who have immunosuppression and HIV infection (regardless of CD4 count) should be given influenza vaccine in accordance with the recommendations and contraindications above. These individuals may not make a full antibody response. Consideration should also be given to the influenza vaccination of household contacts of immunocompromised individuals, i.e. individuals who expect to share living accommodation on most days over the winter and therefore for whom continuing close contact is unavoidable.

There is a theoretical potential for transmission of live attenuated influenza virus in Fluenz Tetra® to immunocompromised contacts for one to two weeks following vaccination. In the US, where there has been extensive use of the live attenuated influenza vaccine, there have been no reported instances of illness or infections from the vaccine virus among immunocompromised patients inadvertently exposed. Where close contact with very severely immunocompromised patients (e.g. bone marrow transplant patients requiring isolation) is likely or unavoidable (for example, household members), however, appropriate alternative inactivated influenza vaccines should be considered.

Further guidance is provided by the Royal College of Paediatrics and Child Health (<http://www.rcpch.ac.uk/>), the British HIV Association

(BHIVA) immunisation guidelines for HIV-infected adults (BHIVA, 2008; <http://www.bhiva.org/immunization-guidelines.aspx>) and the Children's HIV Association (CHIVA) immunisation guidelines (<http://www.chiva.org.uk/professionals/health/guidelines/index.html>).

## Severe asthma or active wheezing

The live attenuated influenza vaccine (LAIV Fluenz Tetra®) is not recommended for children and adolescents with severe asthma or active wheezing, for example those who are currently taking oral steroids or who have been prescribed oral steroids in the last 14 days for respiratory disease. There is limited safety data on children who are currently taking a high dose of an inhaled steroid – Budesonide >800 mcg/day or equivalent (e.g. Fluticasone >500 mcgs/day) – such children should only be given LAIV on the advice of their specialist. As these children are a defined risk group for influenza, those who cannot receive LAIV should receive an inactivated influenza vaccine.

Vaccination with Fluenz Tetra® should be deferred in children with a history of active wheezing in the past 72 hours or those who have increased their use of bronchodilators in the previous 72 hours. If their condition has not improved after a further 72 hours then, to avoid delaying protection in this high risk group, these children should be offered an inactivated influenza vaccine.

Children in this clinical risk group and aged under nine years who have not been previously vaccinated against influenza will require a second dose whether given LAIV or inactivated vaccine.

## Egg allergy

In all settings providing vaccination, facilities should be available and staff trained to recognise and treat anaphylaxis (see Chapter 8).

Inactivated influenza vaccines that are egg-free or have a very low ovalbumin content (<0.12 µg/ml - equivalent to <0.06 µg for a 0.5 ml dose) are available (see below) and studies show they may be used safely in individuals with egg allergy (des Roches et al., 2012). LAIV Fluenz Tetra®, which has an upper ovalbumin limit of 1.2 µg/ml, has also been recently shown (JCVI, 2015) to be safe for use in most egg allergic children (see below). The ovalbumin content of influenza vaccines will be published prior to the influenza season in May of each year in a special edition of vaccine update (<https://www.gov.uk/government/collections/vaccine-update>.)

## Children

JCVI has advised (JCVI, 2015) that, except for those with severe anaphylaxis to egg which has previously required intensive care, children with an egg allergy can be safely vaccinated with Fluenz Tetra® in any setting (including primary care and schools); those with clinical risk factors that contraindicate Fluenz Tetra® should be offered an inactivated influenza vaccine with a very low ovalbumin content (less than 0.12 µg/ml).

Children with a history of severe anaphylaxis to egg which has previously required intensive care, should be referred to specialists for immunisation in hospital. LAIV is not otherwise contraindicated in children with egg allergy. Egg-allergic children with asthma can receive LAIV if their asthma is well-controlled (see the above section on severe asthma).

Children in a clinical risk group and aged under nine years who have not been previously vaccinated against influenza will require a second dose whether given LAIV or inactivated vaccine.

## Adults

The ovalbumin-free influenza vaccine Optaflu®, if available, can be used in any setting on patients from the age of 18 years, regardless of the severity of the egg allergy.

Adult patients can also be immunised in any setting using an inactivated influenza vaccine with an ovalbumin content less than 0.12 µg/ml (equivalent to 0.06 µg for 0.5 ml dose), excepting those with severe anaphylaxis to egg which has previously required intensive care who should be referred to specialists for immunisation in hospital.

## Use with antiviral agents against influenza

There is a potential for influenza antiviral agents to lower the effectiveness of the live attenuated influenza vaccine (Fluenz Tetra®). Therefore, influenza antiviral agents and Fluenz Tetra® should not be administered concomitantly. Fluenz Tetra® should be delayed until 48 hours following the cessation of treatment with influenza antiviral agents. Administration of influenza antiviral agents within two weeks of administration of Fluenz Tetra® may adversely affect the effectiveness of the vaccine.

## Exposure of healthcare professionals to live attenuated influenza vaccine viruses

In theory, healthcare workers may have low level exposure to live attenuated influenza vaccine viruses during administration of the vaccine and/or from

recently vaccinated patients. The vaccine viruses are cold-adapted and attenuated, and are unlikely to cause symptomatic influenza. In the US, where there has been extensive use of the live attenuated influenza vaccine, no transmission of vaccine virus in healthcare settings ever has been reported and there have been no reported instances of illness or infections from the vaccine virus among healthcare professionals inadvertently exposed. Thus, the Centers for Disease Control and Prevention has considered that the risk of acquiring vaccine viruses from the environment is unknown but is probably low (CDC, 2013). As a precaution, however, very severely immunosuppressed individuals should not administer live attenuated influenza vaccine. Other healthcare workers who have less severe immunosuppression or are pregnant, should follow normal clinical practice to avoid inhaling the vaccine and ensure that they themselves are appropriately vaccinated.

### Inadvertent administration of Fluenz Tetra®

If an immunocompromised individual receives LAIV then the degree of immunosuppression should be assessed. If the patient is severely immunocompromised, antiviral prophylaxis should be considered, otherwise they should be advised to seek medical advice if they develop flu-like symptoms in the four days (the usual incubation period) following administration of the vaccine. If antivirals are used for prophylaxis or treatment, then in order to maximise their protection in the forthcoming flu season, the patient should also be offered inactivated influenza vaccine. This can be given straight away.

### Adverse reactions

Pain, swelling or redness at the injection site, low grade fever, malaise, shivering, fatigue, headache, myalgia and arthralgia are among the commonly reported symptoms after intramuscular or intradermal vaccination. A small painless nodule (induration) may also form at the injection site. These symptoms usually disappear within one to two days without treatment. Nasal congestion/rhinorrhoea, reduced appetite, weakness and headache are common adverse reaction following administration of the live attenuated intranasal vaccine (Fluenz Tetra®).

Immediate reactions such as urticaria, angio-oedema, bronchospasm and anaphylaxis can occur.

The following adverse events have been reported very rarely after influenza vaccination over the past 30 years but no causal association has been established: neuralgia, paraesthesiae, convulsions (see note below) and

transient thrombocytopenia, vasculitis with transient renal involvement and neurological disorders such as encephalomyelitis.

A study in the UK found that there was no association between Guillain-Barré syndrome (GBS) and influenza vaccines although there was a strong association between GBS and influenza-like illness. The increased risk of GBS after influenza-like illness, if specific to infection with influenza virus, together with the absence of a causal association with influenza vaccine suggests that influenza vaccine should protect against GBS (Stowe *et al.*, 2009). GBS has been reported very rarely after immunisation with influenza vaccine, one case per million people vaccinated in one US study (Laskey *et al.*, 1998). However, this has not been found in other studies and a causal relationship has not been established (Hurwitz *et al.*, 1981; Kaplan *et al.*, 1982; Roscelli *et al.*, 1991).

Side effects and adverse reactions associated with the influenza vaccines Viroflu<sup>®</sup> and Pandemrix<sup>®</sup> have been previously documented. Viroflu<sup>®</sup> (Janssen-Cilag Ltd, formerly Crucell) may be associated with a higher than expected rate of fever in children aged under five years. An increased risk of narcolepsy after vaccination with the AS03 adjuvanted pandemic A/H1N1 2009 vaccine Pandemrix<sup>®</sup> was identified in England (Miller *et al.*, 2013) consistent with findings first identified in Finland and Sweden (Nohynek *et al.*, 2010; Partinen *et al.*, 2010). Viroflu<sup>®</sup> and Pandemrix<sup>®</sup> are no longer used in the UK influenza immunisation programme

All serious suspected reactions following influenza vaccines should be reported to the Medicines and Healthcare products Regulatory Agency using the Yellow Card scheme at <http://yellowcard.mhra.gov.uk/>

The influenza vaccine Fluenz Tetra<sup>®</sup> and quadrivalent Fluarix<sup>™</sup> Tetra carry a black triangle symbol (▼). This is a standard symbol added to the product information of a vaccine during the earlier stages of its introduction, to encourage reporting of all suspected adverse reactions.

### Febrile convulsions and fever

One inactivated influenza vaccine (Fluvax by bioCSL marketed in the UK by Pfizer as Enzira<sup>®</sup> or influenza vaccine<sup>®</sup>) has been associated with a high rate of febrile convulsions in children under five years of age in other countries. The SPC for Enzira<sup>®</sup> also indicates that a high rate of fever was reported in the age group aged five to under nine years. Due to the risk of febrile convulsions, the indication for Enzira is restricted to use in adults and children aged five years and older. Because of this, this vaccine will not be part of the central supply for use in children in the 2014/15 season, but may be available for purchase by the practice. If no suitable alternative vaccines are available, clinicians should

ensure parents are aware of the risk and give advice on the management of vaccine-induced fever (see **Chapter 8**).

There remains no evidence that other trivalent influenza vaccines used in the UK are associated with a similar risk of febrile convulsions in children (Stowe *et al.*, 2011; Bryan and Seabroke, 2011).

### Management of suspected cases, contacts and outbreaks

There are antiviral drugs available that can be used under certain circumstances to either prevent or treat influenza. NICE has issued guidance on the use of antiviral drugs for the prevention and treatment of influenza at:

<http://publications.nice.org.uk/oseltamivir-amantadine-review-and-zanamivir-for-the-prophylaxis-of-influenza-ta158>

<http://guidance.nice.org.uk/TA168>

It is always important to encourage and maintain good hand and respiratory hygiene which can help to reduce the spread of influenza. Information and resources on the 'Catch it, Bin it, Kill it', hand and respiratory hygiene campaign can be found at:

<https://www.gov.uk/government/organisations/public-health-england>

### Supplies

Demand for influenza vaccine sometimes increases unpredictably in response to speculation about influenza illness in the community. Therefore, it is recommended that practices order sufficient vaccine for their needs, based on their 'at risk' registers, well in advance of the immunisation season.

Information on supplies and how to order vaccines will be given in guidance provided by each of the four UK countries' health departments – see respective websites for details. GPs should order vaccine for those aged 65 years and older and those in adult clinical risk groups from the influenza vaccine manufacturers (contact details below) as in previous years. Live attenuated influenza vaccine (Fluenz Tetra<sup>®</sup>) has been purchased centrally for children aged two to less than 17 years and for children aged 2 years to less than 18 years in risk groups. For children under 18 years of age where Fluenz is medically contraindicated an inactivated trivalent vaccine or Fluarix<sup>™</sup> Tetra will be supplied. These vaccines should be ordered as per the usual mechanisms for the routine childhood immunisation programme (also see **Chapter 3**). Arrangements for supply may differ between England and the devolved administrations.

## Suppliers of influenza vaccines

BGP Products Ltd, formerly Abbott Healthcare 0800 358 7468  
 AstraZeneca UK Ltd 0845 139 0000  
 GlaxoSmithKline 0800 221 441  
 MASTA 0113 238 7552  
 Novartis Vaccines 08457 451 500  
 Pfizer Vaccines 0800 089 4033  
 Sanofi Pasteur MSD 0800 085 5511

A list of the influenza vaccines available in the UK is published ahead of the influenza season in the annual flu letter for England (available at: <https://www.gov.uk/government/collections/annual-flu-programme>).

## References

Allison MA Daley MF, Crane LA *et al.* (2006) Influenza vaccine effectiveness in healthy 6- to 21-month-old children during the 2003-2004 season. *J Pediatr* **149**: 755-62.

Ambrose CS, Levin MJ, Belshe RB. The relative efficacy of trivalent live attenuated and inactivated vaccines in children and adults. *Influenza And Other Respiratory Viruses* 2011;5:67-75.

American Academy of Pediatrics (2003) Active immunization. In: Pickering LK (ed.) *Red Book: 2003 Report of the Committee on Infectious Diseases*, 26th edition. Elk Grove Village, IL: American Academy of Pediatrics, p. 33.

Ampofo K, Gesteland PH, Bender J *et al.* (2006) Epidemiology, complications, and cost of hospitalization in children with laboratory-confirmed influenza infection. *Pediatrics* **118**(6):2409-17.

Ashkenazi S, Vertruyen A, Aristegui J. *et al.* (2006) Superior relative efficacy of live attenuated influenza vaccine compared with inactivated influenza vaccine in young children with recurrent respiratory tract infections. *Pediatr Infect Dis J* 25(10): 870-9. <http://www.ncbi.nlm.nih.gov/sites/entrez/17006279>

Ashkenazi S, Vertruyen A., Aristegui, J. *et al.* [2006] Superior relative efficacy of live attenuated influenza vaccine compared with inactivated influenza vaccine in young children with recurrent respiratory tract infections. *Pediatr Infect Dis J* 870-9.

Barker WH and Mullooly JP (1982) Pneumonia and influenza deaths during epidemics. *Arch Int Med* **142**: 85-9.

Belshe RB, Edwards KM, Vesikari T *et al.* (2007) Live attenuated versus inactivated influenza vaccine in infants and young children. *N Engl J Med* 356(7): 685-96. <http://www.ncbi.nlm.nih.gov/sites/entrez/17301299>

Benowitz I, Esposito DB, Gracey KD *et al.* (2010) Influenza vaccine given to pregnant women reduces hospitalization due to influenza in their infants. *Clin Infect Dis*. 51: 1355-61.

Block S L, Toback SL, Yi T *et al.* (2009) Efficacy of a single dose of live attenuated influenza vaccine in previously unvaccinated children: a post hoc analysis of three studies of children aged 2 to 6 years. *Clin Ther.* **31**: 2140-7.

Bracco Neto H, Farhat CK, Tregnaghi MW, *et al.* (2009) Efficacy and safety of 1 and 2 doses of live attenuated influenza vaccine in vaccine-naive children. *Pediatr Infect Dis J.* **28**: 365-71

British HIV Association (2008) Immunisation guidelines for HIV-infected adults: [www.bhiva.org/documents/Guidelines/Immunisation/Immunization2008.pdf](http://www.bhiva.org/documents/Guidelines/Immunisation/Immunization2008.pdf)  
Accessed: May 2010

Bryan P and Seabroke S (2011) No increased risk of febrile convulsions after seasonal influenza immunisation in UK. *Lancet* **377**: 904.

Centers for Disease Control and Prevention. Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices — United States, 2013–2014 September 20, 2013 / 62(RR07);1- 43 [http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6207a1.htm?s\\_cid=rr6207a1\\_w#LiveAttenuatedInfluenzaVaccines](http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6207a1.htm?s_cid=rr6207a1_w#LiveAttenuatedInfluenzaVaccineshttp://www.cdc.gov/mmwr/preview/mmwrhtml/rr6207a1.htm?s_cid=rr6207a1_w#LiveAttenuatedInfluenzaVaccines)  
[http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6207a1.htm?s\\_cid=rr6207a1\\_w#LiveAttenuatedInfluenzaVaccines](http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6207a1.htm?s_cid=rr6207a1_w#LiveAttenuatedInfluenzaVaccines)

Centers for Disease Control and Prevention (2012) Persons Who Should Not Be Vaccinated Influenza Prevention and Control Recommendations. <http://www.cdc.gov/professionals/acip/shouldnot.htm>

Coffin SE, Zaoutis TE, Rosenquist AB *et al.* (2007) Incidence, complications, and risk factors for prolonged stay in children hospitalized with community-acquired influenza. *Pediatrics* **119**(4):740-8.

Committee on Safety of Medicines (2003) Further data support safety of thiomersal in vaccines. Available from:

[www.mhra.gov.uk/Safetyinformation/Generalsafetyinformationandadvice/Product-specificinformationandadvice/accinesafetyThiomersal\(ethylmercury\)containingvaccines/index.htm](http://www.mhra.gov.uk/Safetyinformation/Generalsafetyinformationandadvice/Product-specificinformationandadvice/accinesafetyThiomersal(ethylmercury)containingvaccines/index.htm)

Accessed: May 2010

Department of Health (2013) Health Technical Memorandum 07-01: Safe management of healthcare waste. [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/167976/HTM\\_07-01\\_Final.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/167976/HTM_07-01_Final.pdf)

Accessed March 2014.

Des Roches A, Paradis L, Gagnon R, *et al.* (2012). Egg-allergic patients can be safely vaccinated against influenza. *J Allergy Clin Immunol.* **130**(5):1213-6.

Dexter LJ, Teare MD, Dexter M *et al.* (2012) Strategies to increase influenza vaccination rates: outcomes of a nationwide cross-sectional survey of UK general practice. *BMJ Open.* **2**:e000851.

Donaldson LJ *et al.* (2010) English mortality from A/H1N1 comparisons with recent flu mortality. *BMJ* **340**: 62.

Eick AA, Uyeki TM, Klimov A *et al.* (2010) Maternal influenza vaccination and effect on influenza virus infection in young infants. *Arch Pediatr Adolesc Med.* **165**: 104-11.

EMA/Committee for Proprietary Medicinal Products (CPMP) Note for guidance on harmonisation of requirement for influenza vaccines. [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Scientific\\_guideline/2009/09/WC500003945.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/09/WC500003945.pdf)  
Accessed March 2014.

EMEA/CPMP/VEG/1194/04 (2004). EMEA public statement on thiomersal in vaccines for human use – recent evidence supports safety of thiomersal-containing vaccines. [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Scientific\\_guideline/2009/09/WC500003904.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/09/WC500003904.pdf)  
Accessed March 2014.

Fezeu L, Julia C, Henegar A, Bitu J *et al.* Obesity is associated with higher risk of intensive care unit admission and death in influenza A (H1N1) patients: a systematic review and meta-analysis *Obes Rev.* 2011 Aug;12(8):653-9.

Fleming DM, Watson JM, Nicholas S *et al.* (1995) Study of the effectiveness of influenza vaccination in the elderly in the epidemic of 1989/90 using a general practice database. *Epidemiol Infect* **115**: 581–9.

Fleming DM, Crovari P, Wahn U *et al.* (2006) Comparison of the efficacy and safety of live attenuated cold-adapted influenza vaccine, trivalent, with trivalent inactivated influenza virus vaccine in children and adolescents with asthma. *Pediatr Infect Dis J* **25**(10): 860-9.  
<http://www.ncbi.nlm.nih.gov/sites/entrez/17006278>

Goddard NL, Kyncl J and Watson JM (2003) Appropriateness of thresholds currently used to describe influenza activity in England. *Commun Dis Public Health* **6**: 238–45.

Heinonen OP, Shapiro S, Monson RR *et al.* (1973) Immunization during pregnancy against poliomyelitis and influenza in relation to childhood malignancy. *Int J Epidemiol* **2**: 229–35.

Heinonen S, Silvennoinen H, Lehtinen P *et al.* (2011) Effectiveness of inactivated influenza vaccine in children aged 9 months to 3 years: an observational cohort study. *Lancet Infect Dis* **11**(1): 23-9.

Hoft DF, Babusis E, Worku S, *et al.* Live and inactivated influenza vaccines induce similar humoral responses, but only live vaccines induce diverse T-cell responses in young children. *J Infect Dis* 2011;204:845–53.

Hurwitz ES, Schonberger LG, Nelson DB *et al.* (1981) Guillain-Barré syndrome and the 1978-1979 influenza vaccine. *N Eng J Med* **304**(26):1557-61.

Jamieson D, Honein MA, Rasmussen SA *et al.* (2009) H1N1 2009 influenza virus infection during pregnancy in the USA. *Lancet* **374**: 451-8.

The Joint Committee for Vaccination and Immunisation statement on the annual influenza vaccination programme – extension of the programme to children 25 July 2012 Available at: [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/224775/JCVI-statement-on-the-annual-influenza-vaccination-programme-25-July-2012.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/224775/JCVI-statement-on-the-annual-influenza-vaccination-programme-25-July-2012.pdf)

Joint Committee on Vaccination and Immunisation Minutes of the February 2015 meeting. Available at : <https://www.gov.uk/government/groups/joint-committee-on-vaccination-and-immunisation#minutes>

Kaplan JE, Katona P, Hurwitz ES *et al.* (1982) Guillain-Barré syndrome in the United States, 1979-80 and 1980-81. *JAMA* **248**(6):698-700

Lasky T, Terracciano GJ, Magder L *et al.* (1998) The Guillain-Barré syndrome and the 1992–1993 and 1993–1994 influenza vaccines. *N Engl J Med* **339**: 1797–1802.

Lau LLH, Cowling BJ, Fang VJ *et al.* (2010) Viral shedding and clinical illness in naturally acquired influenza virus infections. *J Infect Dis* **201**: 1509-16.

Mangtani P, Cumberland P, Hodgson CR *et al.* (2004) A cohort study of the effectiveness of influenza vaccine in older people, performed using the United Kingdom general practice research database. *J Infect Dis* **190**(1): 1–10

McNeil SA, Dodds, LA, Fell, DB *et al.* Effect of respiratory hospitalization during pregnancy on infant outcomes. *Am J Obstet Gynecol* 204: [6 Suppl 1.] S54-7, Epub 2011 Apr 24. 2011 Jun.

Mendez-Figueroa H, Raker, C and Anderson, BL. Neonatal characteristics and outcomes of pregnancies complicated by influenza infection during the 2009 pandemic. *Am J Obstet Gynecol* 204. 204: [6 Suppl 1.] S58-63, Epub 2011 Mar 31. 2011 Jun.

Miller E, Andrews N, Stellitano L, *et al.* Risk of narcolepsy in children and young people receiving AS03 adjuvanted pandemic A/H1N1 2009 influenza vaccine: retrospective analysis. *BMJ* 2013;346: f794.

Morgan OW, Bramley A, Fowlkes A, *et al.* Morbid obesity as a risk factor for hospitalization and death due to 2009 pandemic influenza A(H1N1) disease *PLoS One*. 2010 Mar 15;5(3):

Neuzil KM, Reed, GW, Mitchel, EF *et al.* Impact of influenza on acute cardiopulmonary hospitalizations in pregnant women. *Am J Epidemiol* 148: [11.] 1094-102. 1998 Dec 1.

Neuzil KM, Jackson LA, Nelson J *et al.* (2006) Immunogenicity and reactogenicity of 1 versus 2 doses of trivalent inactivated influenza vaccine in vaccine-naive 5-8-year-old children. *J Infect Dis* **194**(8): 1032-9. <http://www.ncbi.nlm.nih.gov/sites/entrez/16991077>

Nicoll A, Ciancio B, Tsoлова S *et al.* (2008) The scientific basis for offering seasonal influenza immunisation to risk groups in Europe. *Euro Surveill* **13**(43). pii: 19018.

Nohynek H, Jokinen J, Partinen M, *et al.* AS03 adjuvanted AH1N1 vaccine associated with an abrupt increase in the incidence of childhood narcolepsy in Finland. *PLoS One* 2012; 7: e33536.

O'Brien MA, Uyeki TM, Shay DK *et al.* (2004) Incidence of outpatient visits and hospitalizations related to influenza in infants and young children. *Pediatrics* **113**(3 Pt 1):585-93.

Omer SB, Goodman, D, Steinhoff, MC *et al.* Maternal influenza immunization and reduced likelihood of prematurity and small for gestational age births: a retrospective cohort study. *PLoS Med*. 8: [5.] e1000441 Epub 2011 May 31. 2011 May.

Osterholm, MT, Kelley, NS, Sommer, A and Belongia, EA (2012) Efficacy and effectiveness of influenza vaccines: a systematic review and meta-analysis. *Lancet Infect Dis* 12. (1.), 36-44.

Partinen M, Saarenpaa-Heikkila O, Ilveskoski I, *et al.* Increased incidence and clinical picture of childhood narcolepsy following the 2009 H1N1 pandemic vaccination campaign in Finland. *PLoS One* 2012; 7: e33723.

- Pebody R *et al.* (2010) Pandemic influenza A (H1N1) 2009 and mortality in the United Kingdom: risk factors for death, April 2009 to March 2010. *Eurosurveillance* **15**(20): 19571.
- Pierce M, Kurinczuk, JJ, Spark, P *et al.* Perinatal outcomes after maternal 2009/H1N1 infection: national cohort study. *BMJ*. 342. 342:d3214. doi: 10.1136/bmj.d3214.: d3214. 2011 Jun 14.
- Poehling KA, Edwards KM, Weinberg GA *et al.* (2006) The underrecognized burden of influenza in young children. *N Engl J Med* **355**(1):31-40.
- Poehling KA, Szilagyi, PG, Staat, MA *et al.* (2011) Impact of maternal immunization on influenza hospitalizations in infants. *Am J Obstet Gynecol* 204: [6 Suppl 1.] S141-8. Epub Feb 23. 2011 Jun.
- Presanis AM, Pebody RG, Paterson BJ *et al.* Changes in severity of 2009 pandemic A/ H1N1 in England: A Bayesian evidence synthesis. *BMJ*. **343**:d5408 doi:10.1136/bmj.d5408. 2011 Sept 8.
- Ritzwoller DP, Bridges CB, Shetterly S *et al.* (2005) Effectiveness of the 2003-2004 influenza vaccine among children 6 months to 8 years of age, with 1 vs 2 doses. *Pediatrics* **116**: 153-9.
- Roscelli JD, Bass JW, Pang L (1991) Guillain-Barré syndrome and influenza vaccination in the US Army 1980-1988. *Am J Epidemiol* **133**(9):952-5.
- Shuler CM, Iwamoto M, Bridges CB, *et al.* (2007) Vaccine effectiveness against medically attended, laboratory-confirmed influenza among children aged 6 to 59 months, 2003- 2004. *Pediatrics* **119**: e587-95.
- Stowe J, Andrews N, Wise L and Miller E (2009) Investigation of the temporal association of Guillain-Barré Syndrome with influenza vaccine and influenza-like illness using the United Kingdom General Practice Research Database. *Am J Epidemiol* **169**(3):382-8.
- Stowe J, Andrews N, Bryan P, *et al.* (2011) Risk of convulsions in children after monovalent H1N1 (2009) and trivalent influenza vaccines: a database study. *Vaccine* 29, 9467-72.
- Subbramanian RA, Basha S, Shata MT, *et al.*. Pandemic and seasonal H1N1 influenza hemagglutinin-specific T cell responses elicited by seasonal influenza vaccination. *Vaccine* 2010;28:8258–67.
- Tamma PD, Ault KA, del Rio C, Steinhoff MC *et al.* (2009) Safety of influenza vaccination during pregnancy. *Am J Obstet Gynecol* **201**(6): 547-52.
- Therapeutic Goods Administration Australian Government (2010) Investigation into febrile reactions in young children following 2010 seasonal trivalent influenza vaccination. <http://www.tga.gov.au/safety/alerts-medicine-seasonal-flu-100702.htm>
- Thompson WW, Price C, Goodson B *et al.* (2007) Early thiomersal exposure and neuropsychological outcomes at 7 to 10 years. *N Eng J Med* **357**:1281-92.
- Thompson WW, Shay DK, Weintraub E *et al.* (2003) Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA* **289**(2):179-86.
- Thompson WW, Shay DK, Weintraub E *et al.* (2004) Influenza-associated hospitalizations in the United States. *JAMA* **292**(11):1333-40.
- Toback SL, Beigi R, Tennis P *et al.* (2012). Maternal outcomes among pregnant women receiving live attenuated influenza vaccine. *Influenza Other Respi Viruses* **6**, 44-51.

Van Kerkhove MD, WHO Working Group for Risk Factors for Severe H1N1pdm Infection. Risk factors for severe outcomes following 2009 influenza A (H1N1) infection: a global pooled analysis. *PLoS Med.* 2011 Jul;8(7):e1001053.

Wilde JA, McMillan JA, Serwint J *et al.* (1999) Effectiveness of influenza vaccine in health care professionals: a randomised trial. *JAMA* **281**: 908–13.

Wright PF, Thompson J, Vaughn WK *et al.* (1977) Trials of influenza A/New Jersey/76 virus vaccine in normal children: an overview of age-related antigenicity and reactogenicity. *J Infect Dis* **136** (suppl): S731–41.

Writing Committee of the WHO Consultation on Clinical Aspects of Pandemic (H1N1) 2009 Influenza. Clinical aspects of pandemic 2009 influenza A (H1N1) virus infection. *N Engl J Med* **362**: 1708-19.

Zaman K, Roy E, Arifeen SE *et al* (2008) Effectiveness of maternal influenza immunisation in mothers and infants. *N Engl J Med* **359**: 1555-64.

Zhou H, Thompson WW, Viboud CG *et al.* (2012) Hospitalizations associated with influenza and respiratory syncytial virus in the United States, 1993-2008. *Clin Infect Dis* **54**: 1427-36.