

ARE YOU PROTECTED AGAINST HEPATITIS A?

Gay and bisexual men are at risk of **hepatitis A**. Have your hepatitis A vaccine and get protected.



Public Health
England

IT
STARTS
WITH
ME

For more information and the symptoms of hepatitis A, visit:
www.nhs.uk/hepa

This booklet was produced by
Terrence Higgins Trust for
HIV PREVENTION ENGLAND,
and distributed by:



Public Health England product code: 7263500

© Terrence Higgins Trust, June 2018. Terrence Higgins Trust is a registered charity in England and Wales (reg. no. 288527) and in Scotland (SC039986). Company reg. no. 1778149. A company limited by guarantee. If you have any questions or comments about this resource, or would like information on the evidence used to produce it, please email feedback@tht.org.uk. The information included in this publication was correct at the time of going to press. We plan to review this publication within two years. For updates or for the latest information, visit tht.org.uk



THE FACTS ABOUT HEPATITIS A.

WHAT IS HEPATITIS A?

Hepatitis A is inflammation of the liver – usually caused by a viral infection. It is mainly spread through contaminated food, poor hand washing or sex.

During sex it is mainly passed on when licking skin, condoms or sex toys which have faeces (poo) on them. This could include licking skin near the anus (rimming), buttocks, groin or penis (giving oral sex after anal sex).

Gay and bisexual men with multiple partners are particularly at risk.

WHAT ARE THE SYMPTOMS?

Symptoms for hepatitis A are rarely life-threatening. They can be so mild that people may not realise they have it. But weeks after infection it can cause:

- mild flu-like symptoms
- diarrhoea
- itchy skin
- jaundice, meaning your skin and the whites of your eyes turn yellow, your urine is dark and your faeces (poo) are pale.
- nausea
- extreme tiredness
- stomach pain

HOW CAN I PROTECT MYSELF?

You can avoid getting hepatitis A by:

- **Being vaccinated against hepatitis A – available at your sexual health clinic**
- Washing your hands after sex (ideally buttocks, groin and penis too).
- Changing condoms between anal and oral sex.

There is a Hepatitis A vaccine available – ask about it at your next sexual health clinic appointment.

- Covering anything which is shared during sex with a fresh condom or fresh latex glove for each new person.
- Using a latex barrier (eg, a condom cut into a square) for rimming.
- Using latex gloves for fingering or fisting.
- Not sharing sex toys.

HOW IS IT TREATED?

There is no specific treatment for hepatitis A. Your body will clear the infection but this can take several months. Once you clear the infection, you're immune and cannot get it again – but you can get other types of hepatitis.

If you think you might have hepatitis A, or have any questions, visit a sexual health clinic.