


## Revised algorithm<sup>1</sup> for assessing pregnant women with a history of travel during pregnancy to areas with high or moderate risk of Zika virus (ZIKV) transmission<sup>2</sup>

Published (v9) 10 January 2018



## Notes

1. This guidance will be updated as more information becomes available. Currently this algorithm applies to women at all stages of pregnancy although infection in early pregnancy is likely to be the greatest risk.
2. Laboratory testing is performed by the [PHE Rare and Imported Pathogens Laboratory](#) (RIPL). Given the overlap of symptoms and endemic areas with other viral and bacterial infections, RIPL will routinely test significantly symptomatic pregnant women (those hospitalised and/or acutely unwell) returning from areas with active ZIKV transmission for dengue, chikungunya and other infections as well as ZIKV.
3. Assessment of pregnant women should be based on a history of travel to countries and territories reporting high or moderate risk of ZIKV transmission.
4. Clinical illness is suggestive of Zika virus disease if a combination of the following symptoms are reported: rash; itching/pruritis; fever; headache; arthralgia/arthritis; myalgia; conjunctivitis; lower back pain; retro-orbital pain.
5. A pregnant women with typical Zika virus symptoms (as above) that began within 2 weeks of sexual contact with a male sexual partner who has recently travelled (within the previous 8 weeks) to an area with moderate or high risk of ZIKV transmission, should be tested regardless of her own travel history, due the possibility of sexual transmission.
6. Appropriate samples are a clotted blood (or serum) and, if indicated, and/or a small volume of urine without preservative. The samples must be submitted with an appropriate RIPL request form. This form **must** clearly state the pregnancy gestation and both the travel history (ie which countries visited and the dates of the outward and return journeys) and the clinical details (ie the patient's symptoms and the date of illness onset). This is so that the appropriate investigations can be performed and their results correctly interpreted. ZIKV testing will be performed using real-time PCR and serology. For more information refer to: [Zika virus: sample testing advice](#)
7. If Zika virus antibodies are not detected in a serum sample collected 4 or more weeks after the last possible travel-associated or sexual exposure, then recent Zika virus infection is highly unlikely.
8. Neonatologists and obstetricians should collaborate prior to delivery to **agree a plan for investigations at birth**. For more information refer to: [Zika virus congenital infection: algorithm and interim guidance for neonatologists and paediatricians](#)
8. This evaluation and follow-up is likely to include repeat fetal ultrasound and blood testing for ZIKV RNA at four weekly intervals, and consideration of fetal MRI. Abnormal fetal findings will prompt appropriate investigation including, for example, submission of booking and current serum samples for toxoplasma, rubella, parvovirus and CMV serology. Amniocentesis may be considered for ZIKV PCR. For women who have not had symptoms, taking and storing a clotted serum sample locally, without immediate testing, is recommended. In the event that there is a later concern about fetal development, this sample will be available for retrospective testing, including detection of Zika virus antibodies. For more information refer to: [Zika virus: sample testing advice](#)
9. This is in line with [WHO guidance](#) on the management of asymptomatic pregnant women who have returned from Zika-affected countries
10. In this context, 'small fetal head' is defined as: Head Circumference more than 2 Standard Deviations below the mean for gestational age, ie below the 2.5<sup>th</sup> centile.
11. Apart from microcephaly and intracranial calcifications, other brain abnormalities that have been reported in association with ZIKV infection are ventriculomegaly, cell migration abnormalities (eg lissencephaly, pachygyria), and arthrogyryposis (congenital contractures) secondary to central or peripheral nervous system involvement.

© Crown copyright 2018

You may re-use this information (excluding logos) free of charge in any format or medium, under the terms of the Open Government Licence v3.0. To view this licence, visit [OGL](#) or email [psi@nationalarchives.gsi.gov.uk](mailto:psi@nationalarchives.gsi.gov.uk). Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned. Published (v9) 10 January 2018. PHE publications gateway number: 2015632